

BREWERS' YEAST-MEDIATED SYNTHESIS OF (1*S*,2*S*)-1-(2-THIENYL)-1,2-PROPANEDIOL AND A STUDY ON THE LIPASE-CATALYZED REGIOSELECTIVE INTRODUCTION OF ACYL PROTECTIVE GROUP TO THE DIOL MOIETY

Naoki Mochizuki,[†] Takeshi Sugai,^{††} Hiromichi Ohta,^{††}
Tsutomu Yokomatsu,[¶] and Shiroshi Shibuya*[¶]

[†]Alcohol Beverage Research and Development Laboratory, Asahi
Breweries Ltd., 2-13-1 Ohmori-kita, Ohta-ku, Tokyo 143, Japan

^{††}Department of Chemistry, Keio University, 3-14-1, Hiyoshi, Yokohama
223, Japan


[¶]School of Pharmacy, Tokyo University of Pharmacy and Life Science,
1432-1 Horinouchi, Hachioji, Tokyo 192-03, Japan

Abstract – Reductive homologation of thiophenecarboxyaldehyde mediated by brewers' yeast and subsequent recrystallization provided (1*S*,2*S*)-1-(2-thienyl)-1,2-propanediol in 29% yield. A *Pseudomonas* lipase-catalyzed acetylation preferentially occurred on the hydroxyl group at C-1 position of the diol.

In our previous paper,¹ we reported a brewers' yeast-mediated reductive bishomologation of furfural (**1**). Along with furfuryl alcohol, (1*S*,2*S*)-1-(2-furyl)-1,2-propanediol (**2**) was obtained in a highly enantioselective manner (Scheme 1).

Recent reports on the biochemical transformation²⁻⁵ of thiophene-containing compounds with a substantial synthetic potential⁶ prompted us to study the preparation of (1*S*,2*S*)-1-(2-thienyl)-1,2-propanediol (**3a**) from thiophenecarboxyaldehyde (**4**).

Scheme 1



The incubation of **4** with brewers' yeast¹ gave a mixture of **3a** (34%) and 2-thienylmethanol (52%). The diastereomeric ratio of **3a** was 97 : 3 as judged from its ¹H-NMR spectrum. The *anti*-isomer was elucidated to be the major product, by the comparison of coupling constants and NOE enhancement of the corresponding acetonides (**5**) and (**7**) with those of **6** and **8**, whose relative configuration had been unambiguously determined¹ (Scheme 2). The results are listed in Table 1. The major *anti*-isomer was obtained as a pure state by recrystallization of the mixture.

Scheme 2






Table 1

	5	6	7	8
J_{ab} (Hz)	6.5	6.3	8.5	8.6
NOE ($H_a \rightarrow H_b$, %)	7.1	7.2	—	—

The next task was the determination of the absolute configuration of the purified isomer. In a similar manner to our previous report,¹ both enantiomers [(1*S*,2*S*) and (1*R*,2*R*) isomer] were prepared from α -siloxyaldehydes ((*S*)-**9** and (*R*)-**9**), respectively. Then, the adducts were converted to the corresponding MTPA esters (**3b**) and analyzed by ¹H-NMR (Scheme 3). The results clearly showed that the absolute configuration of major *anti*-isomer of **3a** was (1*S*,2*S*) and the recrystallized product turned out to be isomerically pure.

Scheme 3



On the other hand, precise determination of the ee and the absolute configuration for *syn*-**3a** was not possible due to some difficulty with the isolation in a pure state. The previous study of brewers' yeast-mediated reductive homologation with benzaldehyde and furfural shows that the e.e.s of minor *syn*-isomers are as low as 9-33% ee.¹ Accordingly the ee of *syn*-**3a** might be estimated to be not high.

The structure of (1*S*,2*S*)-**3a** including vicinal glycol system is well coincided with the partial structure of L-rhamnose, a useful optically active starting material in synthetic organic chemistry, in its carbon skeleton as well as the stereochemical sense.⁷ As pure (1*S*,2*S*)-**3a** became in hand, we then attempted a selective introduction of a protective group on the vicinal glycol system, since a regioselectively protected form would increase its synthetic utility for a chiral synthon. The differentiation of two secondary hydroxyl groups turned out to be difficult by a conventional chemical reaction; for example, benzylation of **3a** with even a limited amount of benzoic anhydride under mild conditions afforded a mixture of monobenzoates and dibenzoate. We realized that the reactivity of the two hydroxyl groups is very similar toward acylating agent and turned our attention to biocatalyst-mediated selective reaction, especially a *Pseudomonas* lipase-catalyzed acetylation.⁸ So far, in many reports, (*S*)-isomers of methylcarbinols have been reported to be unreactive.⁹ These characteristics would enable the preferential acetylation of the hydroxyl group on C-1 position, because of the possibly lower reactivity of the hydroxyl group on C-2 position of (1*S*,2*S*)-**3a**.

Scheme 4




Table 2

substrate	wt of lipase / wt of substrate	reaction time	total Yd of mono- acetates (%)	1-acetate (3c) : 2-acetate (3d)
(1 <i>S</i> ,2 <i>S</i>)- 3a	2.0	5 d	35	4.1 : 1
(1 <i>R</i> ,2 <i>R</i>)- 3a	0.5	30 h	80	1 : 5.0

Although the reaction proceeded rather slowly, a preferential acetylation of the hydroxyl group at C-1 position of (1*S*,2*S*)-**3a** took place as expected on treatment with the lipase in vinyl acetate. The ratio of the products (total 35%), 1-acetate (**3c**) to 2-acetate (**3d**) was 4.1 : 1, as judged by ¹H-NMR measurement. At this stage, we became interested in the lipase-catalyzed acetylation of the antipodal isomer, which would be useful to examine the stereochemistry-regioselectivity relationship of the reaction.

In contrast to the previous case, when (1*R*,2*R*)-**3a** was used as the substrate, the reaction was very fast and a preferential acetylation of the hydroxyl group at C-2 position (1-acetate (**3c**) : 2-acetate (**3d**) = 1 : 5) of the product (80%) was observed. The both results are summarized in Table 2. The order of reactivity of hydroxyl groups was concluded to be [(2*R*)-OH > (1*R*)-OH > (1*S*)-OH > (2*S*)-OH]. The highest reactivity of (2*R*)-OH of **3a** among four secondary alcohols well coincides with a recently reported result, which had been observed in the case of the lipase-catalyzed regioselective hydrolysis of the acetyl group of ethyl 3-aryl-2,3-diacetoxypropanoates.¹⁰ The decreased regioselectivity and yield of the transesterification on (1*S*,2*S*)-**3a** might also reflect on the bulkyness of the heterocyclic ring in steric hindrance.^{10,11}

In conclusion, we have established a short synthesis of enantiomerically and diastereomerically pure (1*S*,2*S*)-1-(2-thienyl)-1,2-propanediol from thiophenecarboxyaldehyde. In addition, the reactivity of the secondary alcohols of the product in the course of *Pseudomonas* lipase-catalyzed acetylation was clarified.

EXPERIMENTAL

Mps were uncorrected. IR spectrum was measured as KBr disc on a Perkin-Elmer 1710 FTIR spectrophotometer. ¹H-NMR spectra were measured in CDCl₃ with TMS as the internal standard at 400 MHz on a Bruker AM 400 or JEOL JNM α-400 spectrometer. Optical rotations were recorded on a Jasco DIP 360 polarimeter. Mass spectra were measured on a Hitachi M-80B or a VG Auto Spec spectrometer by the EI method.

(1*S*,2*S*)-1-(2-Thienyl)-1,2-Propanediol (3a)

The incubation condition was followed by the previous report.¹ Starting from 4.4 g of **4**, **3a** (2.1 g, 34%) was obtained as an oil; a mixture of *anti* and *syn* isomers in a 97 : 3 ratio. ¹H-NMR (400 MHz, CDCl₃) δ: 1.13 (*syn*, d, 3H, *J* = 6.4 Hz), 1.15 (*anti*, d, 3H, *J* = 6.4 Hz), 3.93 (*syn*, dq, 1H, *J* = 7.1, 6.4 Hz), 4.07 (*anti*, dq, 1H, *J* = 4.0, 6.4 Hz), 4.41 (*syn*, d, 1H, *J* = 7.1 Hz), 4.87 (*anti*, d, 1H, *J* = 4.0 Hz).

The corresponding acetonides were prepared in a conventional manner: major *cis*-acetonide (**5**) (from *anti*-**3a**) and minor *trans*-acetonide (**7**) (from *syn*-**3a**) were obtained as an inseparable mixture. Measurement of the NMR spectra and a comparison with the spectra of the corresponding acetonides (**6** and **8**) from **2**¹ revealed the relative stereochemistry of **5** and **7**. ¹H-NMR (400 MHz, CDCl₃) δ: 0.99 (**5**, d, 3H, *J* = 6.4 Hz), 1.32 (**7**, d, 3H, *J* = 6.0 Hz), 1.44 (**5**, s, 3H), 1.50 (**7**, s, 3H), 1.53 (**7**, s, 3H), 1.64 (**5**, s, 3H), 4.02 (**7**, H-5, dq, 1H, *J* = 6.0, 8.5 Hz), 4.53 (**5**, H-5, dq, 1H, *J* = 6.4, 6.5 Hz), 4.73 (**7**, H-4, 1H, d, *J* = 8.5 Hz), 5.40 (**5**, H-4, d, 1H, *J* = 6.5 Hz), 6.91 (**5**, broad d, 1H, *J* = 3.5 Hz), 6.98 (**5**, dd, 1H, *J* = 3.5, 5.1 Hz), 7.04 (**7**, 1H, broad d, *J* = 2.5 Hz), 7.26 (**5**, dd, 1H, *J* = 1.2, 5.1 Hz), 7.30 (**7**, dd, 1H, *J* = 1.1, 4.9 Hz). MS *m/z* (rel. int.): 198 (M⁺, 5%), 183 (M⁺-Me, 8%), 154 (M⁺-C₂H₄O, 84%), 141 (25%), 125 (50%), 96 (100%), 58 (42%), 43 (64%). It has been reported that *J*_{4,5} = 8.6 Hz for the *trans*-acetonide (**8**) and *J*_{4,5} = 6.3 Hz for the *cis*-acetonide (**6**).¹ Irradiation at H-4 in **5** (δ 5.40) enhanced the signal of H-5 (δ

4.53) by 7.1%, while no such enhancement was observed in the case of irradiating H-4 (δ 4.73) in **7**.

Diol (**3a**) solidified in a refrigerator, and recrystallization of the mixture from diisopropyl ether afforded pure *anti*-**3a** as needles, mp 59-60°C in 85% recovery, Anal. Calcd for C₇H₁₀O₂S: C, 53.14; H, 6.37. Found: C, 53.17; H, 6.40. $[\alpha]_D^{28} -15.6^\circ$ (*c* 1.36, CHCl₃). IR (cm⁻¹): 3270, 1443, 1372, 1346, 1322, 1277, 1235, 1131, 1081, 1066, 992, 924. MS *m/z* (rel. int.): 158 (M⁺, 4%), 141 (M⁺-OH, 26%), 113 (C₅H₅OS⁺, 89%), 97 (48%), 85 (100%), 81 (37%), 45 (84%). NMR measurements indicated that the crystal of **3a** was diastereomerically pure; ¹H-NMR (400 MHz, CDCl₃) δ : 1.15 (d, 3H, *J* = 6.4 Hz), 4.07 (dq, 1H, *J* = 4.0, 6.4 Hz), 4.87 (d, 1H, *J* = 4.0 Hz), 7.00 (dd, 1H, *J* = 3.5, 4.8 Hz) 7.03 (dd, 1H, *J* = 1.3, 3.5 Hz), 7.31 (dd, 1H, *J* = 1.3, 4.8 Hz).

Determination of absolute configuration

In a similar manner as described before,¹ a diastereomeric mixture of **3a** (*anti* : *syn* = 60 : 1) was prepared from aldehyde ((*S*)-**9**) and was subsequently recrystallized to give an authentic sample: mp 59-60°C; $[\alpha]_D^{28} -15.6^\circ$ (*c* 1.06, CHCl₃). This was converted to the corresponding (*R*)-MTPA esters ((1*S*,2*S*)-**3b**). In the same manner, (1*R*,2*R*)-**3b** was prepared from (*R*)-**9** via (1*R*,2*R*)-**3a**: mp 59-60°C; $[\alpha]_D^{28} +15.6^\circ$ (*c* 1.05, CHCl₃). ¹H-NMR (400 MHz, CDCl₃) δ (1*S*,2*S*)-**3b**: 1.35 (d, 3H, *J* = 6.5 Hz), 3.42 (s, 3H), 3.44 (s, 3H), 5.56 (dq, 1H, *J* = 4.2, 6.5 Hz), 6.04 (d, 1H, *J* = 4.2 Hz), 6.91 (dd, 1H, *J* = 4.0, 5.0 Hz), 6.96 (d, 1H, *J* = 4.0 Hz), 7.25-7.44 (m, 11H); (1*R*,2*R*)-**3b**: 1.17 (d, 3H, *J* = 6.4 Hz), 3.28 (s, 3H), 3.38 (s, 3H), 5.50 (dq, 1H, *J* = 5.5, 6.4 Hz), 6.33 (d, 1H, *J* = 5.5 Hz), 6.98 (dd, 1H, *J* = 3.6, 5.0 Hz), 7.17 (d, 1H, *J* = 3.6 Hz), 7.26-7.34 (m, 11H). By comparing its NMR spectrum with that of an authentic specimen, the sample of **3b** from **3a** obtained by recrystallization of the yeast-fermentation product was determined to be a (1*S*,2*S*)-isomer, and was diastereomerically and enantiomerically pure.

Regioselective acetylation with *Pseudomonas* lipase PS

A mixture of (1*S*,2*S*)-**3a** (20.8 mg), *Pseudomonas* lipase (Amano PS, 40 mg) and vinyl acetate (200 μ L) was stirred at 30°C for 5 days. The mixture was filtered with a pad of Celite and the solid residue was washed with ethyl acetate. The filtrate and washings were combined and concentrated in vacuo, and the residue was purified by a preparative thin-layer chromatography (Merck 5744, 20 cm x 10 cm) developed with hexane-ethyl acetate (1 : 1). A mixture of 1-

acetate and 2-acetate in a 4.1 : 1 ratio was obtained (9.3 mg, 35%). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 1.18 (1-acetate, d, 3H, $J = 6.4$ Hz), 1.22 (2-acetate, d, 3H, $J = 6.4$ Hz), 2.08 (2-acetate, s, 3H), 2.10 (1-acetate, s, 3H), 2.51 (2-acetate, d, 1H, OH, $J = 3.9$ Hz), 4.15 (1-acetate, dq, 1H, $J = 4.9, 6.4$ Hz), 5.05 (2-acetate, ddd, 1H, $J = 1.3, 3.9, 3.9$ Hz), 5.19 (2-acetate, dq, 1H, $J = 3.9, 6.4$ Hz), 5.93 (1-acetate, d, 1H, $J = 4.9$ Hz), 6.99 (2-acetate, dd, 1H, $J = 3.4, 4.9$ Hz), 7.00 (1-acetate, dd, 1H, $J = 3.7, 5.1$ Hz), 7.02 (2-acetate, ddd, 1H, $J = 1.3, 1.3, 3.4$ Hz), 7.14 (1-acetate, dd, 1H, $J = 1.2, 3.7$ Hz), 7.28 (2-acetate, dd, 1H, $J = 1.3, 4.9$ Hz), 7.32 (1-acetate, dd, 1H, $J = 1.2, 5.1$ Hz). MS m/z (rel. int.): 156 ($\text{M}^+ - \text{COCH}_3$, 19%), 140 ($\text{M}^+ - \text{CH}_3\text{CO}_2\text{H}$, 38%), 113 ($\text{C}_5\text{H}_5\text{OS}^+$, 100%), 97 (19%), 85 (32%), 43 (91%).

In a similar manner, (1*R*,2*R*)-**3a** (20.2 mg) was treated with Amano PS (10 mg) in vinyl acetate (200 μL) at 30°C for 30 h. Subsequent workup and purification afforded a mixture of 1-acetate and 2-acetate in a 1 : 5 ratio (20.0 mg, 80%).

ACKNOWLEDGMENTS

The authors thank Amano Pharmaceutical Co. for generous gift of lipase PS and Dr. Yukio Hitotsuyanagi for his technical assistance on the NOE experiment.

REFERENCES AND NOTE

1. N. Mochizuki, S. Hiramatsu, T. Sugai, H. Ohta, H. Morita, and H. Itokawa, *Biosci. Biotech. Biochem.*, 1995, **59**, 2282.
2. K. Miyamoto and H. Ohta, *Biocatalysis*, 1991, **5**, 49.
3. B. M. Kim, H.-Y. Lee, P. M. Munson, J. P. Guare, and C. McDonough, *Tetrahedron Lett.*, 1993, **34**, 6517.
4. F. Bracher and T. Papke, *Tetrahedron: Asymmetry*, 1994, **5**, 1653.
5. S.-K. Kang, J.-H. Jeon, T. Yamaguchi, J.-S. Kim, and B.-S. Ko, *Tetrahedron: Asymmetry*, 1995, **6**, 2139.
6. S. Gronowitz, *Advances in Heterocyclic Chemistry: Recent Advances in the Chemistry of Thiophenes*, Vol. 1, ed. by A. R. Katritzky, Academic Press, Inc., London, 1963, pp. 1-124; G. L'abbé, *Tetrahedron*, 1982, **38**, 3537; A. R. Katritzky and C. W. Rees, *Comprehensive*

- Heterocyclic Chemistry: the Structure, Reactions, Synthesis and Uses of Heterocyclic Compounds, Vol 4, Part 3, Pergamon Press Ltd., Oxford, 1984; S. Gronowitz, The Chemistry of Heterocyclic Compounds: Thiophene and Its Derivatives, Vol 44, Part 1, John Wiley & Sons, Inc., New York, 1985; G. Cardillo and M. Orena, *Tetrahedron*, 1990, **46**, 3321.
7. E. C. Taylor, P. A. Jacobi, *J. Am. Chem. Soc.*, 1974, **96**, 6781.
 8. T. Yokomatsu, N. Nakabayashi, K. Matsumoto, and S. Shibuya, *Tetrahedron: Asymmetry*, 1995, **6**, 3055; A. Sobti and G. A. Sulikowski, *Tetrahedron Lett.*, 1995, **36**, 4193; M.-J. Kim, G.-B. Choi, J.-Y. Kim, and H.-J. Kim, *Tetrahedron Lett.*, 1995, **36**, 6253; K. Matsumoto, S. Fuwa, and H. Kitajima, *Tetrahedron Lett.*, 1995, **36**, 6499; E. Mizuguchi, T. Suzuki, and K. Achiwa, *Synlett*, 1996, 743; T. Sugai, H. Ikeda, and H. Ohta, *Tetrahedron*, 1996, **52**, 8123; K. Lemke, F. Theil, A. Kunath, and H. Schick, *Tetrahedron: Asymmetry*, 1996, **4**, 971; T. Itoh, A. Uzu, N. Kanda, and Y. Takagi, *Tetrahedron Lett.*, 1996, **37**, 91; O. Houille, T. Schmittberger, and D. Uguen, *Tetrahedron Lett.*, 1996, **37**, 625; T. Itoh, M. Shiromoto, H. Inoue, H. Hamada, and K. Nakamura, *Tetrahedron Lett.*, 1996, **37**, 5001; T. Yokomatsu, M. Sato, and S. Shibuya, *Tetrahedron: Asymmetry*, 1996, **7**, 2743.
 9. K. Burgess and L. D. Jennings, *J. Am. Chem. Soc.*, 1991, **113**, 6129; R. J. Kazlauskas, A. N. E. Weissfoch, A. T. Rappaport, and L. A. Cuccia, *J. Org. Chem.*, 1991, **56**, 2656; K. Nakamura, M. Kawasaki, and A. Ohno, *Bull. Chem. Soc. Jpn.*, 1996, **69**, 1079; H. Hamada, M. Shiromoto, M. Funahashi, T. Itoh, and K. Nakamura, *J. Org. Chem.*, 1996, **61**, 2332; M. Kinoshita and A. Ohno, *Tetrahedron*, 1996, **52**, 5397.
 10. S. B. Desai, N. P. Argade, and K. N. Ganesh, *J. Org. Chem.*, 1996, **61**, 6730; As another example for the differentiation of vicinal diols by a chemical method, see T. Yokomatsu, K. Suemune, T. Yamagishi, and S. Shibuya, *Synlett*, 1995, 847.
 11. D. G. Drueckhammer, C. F. Barbas III, K. Nozaki, C.-H. Wong, C. Y. Wood, and M. A. Ciufolini, *J. Org. Chem.*, 1988, **53**, 1607; P. Allevi, P. Ciuffreda, G. Tarocco, and M. Anastasia, *J. Org. Chem.*, 1996, **61**, 4144; J. Kaminska, I. Górnicka, M. Sikora, and J. Góra, *Tetrahedron: Asymmetry*, 1996, **7**, 907.