


THE SYNTHESIS OF 5'-ALKYLNICOTINES

Makoto Shibagaki, Hajime Matsushita, and Hajime Kaneko
 Central Research Institute, Japan Tobacco INC.,
 6-2 Umegaoka, Midori-ku, Yokohama, Kanagawa 227, Japan

Abstract - The reaction of cotinine(2) with alkyl lithium followed by the reduction with NaBH₃CN in acidic methanol gave *cis* and *trans*-5'-alkylnicotines(3 and 4). The stereochemistry of the products was discussed.

To understand the biological activities of nicotine(1), the synthesis of its analogues for the development of structure-activity relationship is important. Although several alkylnicotines have been prepared from this point of view,¹ as to 5'-alkylnicotines only 5'-dimethylnicotine has so far been obtained.² In this paper, we wish to report the synthesis of 5'-monoalkylnicotines. The stereochemistry will also be mentioned.

The reaction of cotinine(2) with alkyl lithium followed by the reduction with sodium cyanoborohydride in acidic methanol gave 5'-monoalkylnicotine. The yield and the stereoselectivity were listed in Table 1. The alkylation of 2



with methyl, ethyl, *n*-butyl, isopropyl and phenyl lithium gave *cis*- and *trans*-5'-alkylnicotines (3 and 4) in moderate yields. With *tert*-butyl lithium, *cis*-isomer only was isolated in very low yield.

The relative configurations of 3 and 4 were confirmed by the NMR analysis. The assignments of individual resonances in these compounds shown in Table 2 and

Table 3 were made by H-H and C-H decoupling methods. As shown in Table 2, the chemical shifts of 5'-proton in 4 were observed at lower field than those in 3. Since 5'- α (pseudo equatorial)-proton in 1 is assigned to the lower field resonance than 5'- β (pseudo axial)-proton,³ 3 and 4 were assumed to have pseudo axial and pseudo equatorial protons, respectively. This explanation was further supported by

¹³C NMR analysis. As shown in Table 3, the chemical shifts of 2'- and 5'-carbons in 3 were observed at lower field than those in 4. This difference was explained to be caused by the steric effect from 5'-substituent. An axial substituent causes higher field shift of α and γ carbons than an equatorial one in cyclohexane ring.⁴ Therefore, the 5'-substituents in 3 and 4 were considered to be pseudo equatorial and pseudo axial, respectively. Two possible conformations of cis-5'-alkyl isomer were anticipated, but the conformation of pyridine ring and alkyl group being both pseudo equatorial was more stable than that being both pseudo axial. Then the compounds possessing 5'-pseudo axial proton and pseudo equatorial substituent, that is 3, were cis-isomers.

As shown in Table 1, cis-isomers were predominantly obtained, and the stereoselectivity increased with an increase in the bulkiness of 5'-alkyl

Table 1 The yield and stereoselectivity

| substituent | yield(%) | <u>3</u> : <u>4</u> (%) |
|----------------------|----------|-------------------------|
| a methyl | 68 | 78 : 22 |
| b ethyl | 47 | 82 : 18 |
| c n-butyl | 70 | 82 : 18 |
| d phenyl | 66 | 90 : 10 |
| e isopropyl | 54 | 90 : 10 |
| f <u>tert</u> -butyl | 9 | 100 : 0 |





Table 2 ^1H NMR chemical shift (ppm from TMS)

| 1^* | $3a$ | $4a$ | $3b$ | $4b$ | $3c$ | $4c$ | $3d$ | $4d$ | $3e$ | $4e$ | $3f$ | |
|-------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1' | 2.18 | 2.11 | 2.16 | 2.11 | 2.16 | 2.11 | 2.15 | 2.01 | 1.92 | 2.10 | 2.15 | 2.31 |
| 2' | 3.07 | 3.24 | 3.72 | 3.24 | 3.80 | 3.23 | 3.80 | 3.44 | 4.14 | 3.31 | 4.14 | 3.51 |
| 3' | 1.73 | 1.66 | 1.76 | 1.60 | 1.76 | 1.61 | 1.76 | 1.81 | 1.96 | 1.53 | 1.78 | 1.59 |
| | 2.21 | 2.10 | 2.37 | 2.08 | 2.32 | 2.08 | 2.30 | 2.23 | 2.54 | 2.05 | 2.48 | 1.97 |
| 4' | 1.80 | 1.57 | 1.58 | 1.60 | 1.57 | 1.56 | 1.67 | 1.81 | 1.96 | 1.67 | 1.52 | 1.77 |
| | 1.95 | 1.98 | 2.24 | 1.96 | 2.12 | 1.95 | 2.14 | 2.23 | 2.54 | 1.72 | 2.27 | 1.92 |
| 5' | 2.31 | 2.42 | 3.41 | 2.31 | 3.02 | 2.33 | 3.07 | 3.44 | 4.14 | 2.39 | 3.29 | 2.43 |
| | 3.25 | - | - | - | - | - | - | - | - | - | - | - |
| 6' | - | 1.21 | 1.06 | 1.38 | 1.21 | 1.32 | 1.22 | - | - | 1.91 | 1.93 | - |
| | - | - | - | 1.74 | 1.95 | 1.71 | 1.65 | - | - | - | - | - |
| 7' | - | - | - | 0.95 | 0.91 | 1.36 | 1.34 | 7.45 | 7.46 | 0.93 | 0.94 | 0.94 |
| | - | - | - | - | - | - | - | - | - | 0.95 | 0.95 | - |
| 8' | - | - | - | - | - | 1.37 | 1.37 | 7.35 | 7.34 | - | - | - |
| 9' | - | - | - | - | - | 0.93 | 0.93 | 7.27 | 7.27 | - | - | - |
| 2 | 8.54 | 8.53 | 8.51 | 8.54 | 8.50 | 8.53 | 8.49 | 8.65 | 8.54 | 8.55 | 8.54 | 8.61 |
| 4 | 7.68 | 7.74 | 7.65 | 7.71 | 7.61 | 7.70 | 7.61 | 7.83 | 7.64 | 7.71 | 7.53 | 7.77 |
| 5 | 7.22 | 7.25 | 7.25 | 7.23 | 7.24 | 7.24 | 7.24 | 7.27 | 7.27 | 7.23 | 7.24 | 7.23 |
| 6 | 8.48 | 8.49 | 8.48 | 8.48 | 8.48 | 8.47 | 8.47 | 8.51 | 8.51 | 8.47 | 8.47 | 8.47 |

* see reference 3

groups; *tert*-butyl > *iso*-propyl = phenyl > methyl = ethyl = *n*-butyl. This reaction is presumed to proceed as shown in Scheme. The reaction of **2** with alkyl lithium gives hydroxy derivative(**5**), which is dehydrated to enamine derivative(**6**) in acidic media.⁵ It is well known that the reduction of enamine by sodium



Figure

cyanoborohydride in acidic media proceeded via iminium salt. Therefore the reduction of **6** proceeds via iminium intermediate(**7**). As shown in Figure, a

Table 3 ^{13}C NMR chemical shift (ppm from TMS)

| 1^* | $3a$ | $4a$ | $3b$ | $4b$ | $3c$ | $4c$ | $3d$ | $4d$ | $3e$ | $4e$ | $3f$ | |
|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1' | 40.3 | 38.5 | 35.1 | 39.0 | 35.2 | 39.0 | 35.2 | 38.7 | 35.5 | 39.1 | 39.2 | 43.9 |
| 2' | 68.8 | 70.0 | 64.5 | 70.0 | 65.4 | 70.0 | 64.8 | 69.1 | 65.7 | 69.8 | 68.9 | 71.6 |
| 3' | 35.2 | 33.2 | 31.9 | 33.8 | 33.0 | 33.6 | 33.1 | 34.2 | 33.5 | 34.6 | 34.7 | 36.3 |
| 4' | 22.7 | 32.0 | 29.9 | 29.2 | 28.9 | 29.7 | 29.4 | 34.1 | 33.3 | 23.7 | 23.7 | 26.9 |
| 5' | 56.9 | 62.4 | 58.6 | 68.2 | 65.1 | 67.1 | 63.6 | 71.7 | 68.1 | 71.2 | 71.2 | 74.7 |
| 6' | - | 19.6 | 14.9 | 27.0 | 22.2 | 34.2 | 29.5 | 144.0 | 143.2 | 29.3 | 29.2 | 36.2 |
| 7' | - | - | - | 10.3 | 11.1 | 28.5 | 29.1 | 127.5 | 128.0 | 15.5 | 15.5 | 27.2 |
| | - | - | - | - | - | - | - | - | - | 20.1 | 20.0 | - |
| 8' | - | - | - | - | - | 23.3 | 23.2 | 128.5 | 128.4 | - | - | - |
| 9' | - | - | - | - | - | 14.3 | 14.3 | 127.2 | 127.2 | - | - | - |
| 2 | 149.4 | 149.7 | 149.8 | 149.6 | 149.9 | 149.7 | 149.8 | 149.6 | 149.8 | 149.6 | 149.6 | 149.5 |
| 3 | 138.6 | 139.8 | 139.8 | 139.9 | 139.9 | 139.8 | 139.8 | 139.8 | 139.8 | 140.1 | 140.1 | 140.1 |
| 4 | 134.6 | 135.2 | 135.2 | 135.0 | 135.3 | 135.1 | 135.3 | 135.0 | 135.3 | 134.9 | 135.4 | 134.6 |
| 5 | 123.3 | 123.7 | 123.6 | 123.6 | 123.5 | 123.7 | 123.6 | 123.7 | 123.5 | 123.6 | 123.2 | 123.5 |
| 6 | 148.4 | 148.7 | 148.6 | 148.5 | 148.6 | 148.7 | 148.6 | 148.6 | 148.7 | 148.5 | 148.3 | 148.4 |

* see reference 3

hydride attacks C=N double bond from side B predominantly to give 3 , because side A is hindered by 2'-pyridine ring.

EXPERIMENTAL

2'S,5'S-5'-Methylnicotine(3a) and 2'S,5'R-5'-Methylnicotine(4a) --- To a solution of cotinine(2) (5mmol, 0.9g) in 20ml of ether was added dropwise 6mmol of methyl lithium in 10ml of ether at 0°C. The reaction mixture was stirred at room temperature. After 1 h, dil. HCl was added to the mixture, made basic with NaHCO_3 , and extracted with ether. The extract was dried over Na_2SO_4 , and concentrated *in vacuo*. The resulting oil was dissolved in 20ml of methanol, and a trace of bromocresol green (pH 3.8-5.4) was added followed by

the addition of a 2N solution of HCl in methanol until the color turned yellow. Then 0.33g(5mmol) of NaBH₃CN was added with stirring, and the methanolic HCl was added dropwise to maintain the yellow color. After stirring at room temperature for 1h, the solution was poured into 20ml of 0.1N NaOH, saturated with NaCl and extracted with ether. The extract was dried over Na₂SO₄, an concentrated *in vacuo* to give a mixture of 2'S,5'S- and 2'S,5'R-5'-methylnicotine(3a and 4a). These diastereomers were separated by a reversed phase column chromatography (ODS; 22mm ϕ x 50cm; CH₃OH : H₂O = 4 : 1), and 0.47g(53%) of 3a and 0.13g(15%) of 4a were obtained. (3a): mass(m/z); 176(M+; 7), 98(22), 118(7), 130(35), 147(6), 161(100), 162(10), 175(5): Found: m/z, 176.1322. Calcd for C₁₁H₁₆N₂: M, 176.1314; [α]₅₄₆²⁵ -96.4° (c=0.6, MeOH); (4a): mass(m/z); 176(M+; 10), 98(18), 118(7), 130(41), 147(12), 161(100), 162(11), 175(8): Found: m/z, 176.1325. Calcd for C₁₁H₁₆N₂: M, 176.1314; [α]₅₄₆²⁵ -65.2° (c=0.23, MeOH).

2'S,5'S-5'-Ethylnicotine(3b) and 2'S,5'R-5'-Ethylnicotine(4b) --- The reaction of 2 with ethyl lithium was carried out in a similar manner as described above. After separation, 0.37g(39%) of 2'S,5'S-5'-ethylnicotine(3b) and 0.08g(8%) of 2'S,5'R-5'-ethylnicotine(4b) were obtained. (3b): mass(m/z); 190(M+; 2), 118(7), 130(23), 161(100), 162(11): Found: m/z, 190.1433. Calcd for C₁₂H₁₈N₂: M, 190.1471; [α]_D²⁵ -40.7° (c=1.15, MeOH); (4b): mass(m/z); 190(M+; 4), 118(7), 130(42), 161(100), 162(15): Found: m/z, 190.1450. Calcd for C₁₂H₁₈N₂: M, 190.1471; [α]_D²⁵ -65.5° (c=0.33, MeOH).

2'S,5'S-5'-n-Butylnicotine(3c) and 2'S,5'R-5'-n-Butylnicotine(4c) --- The reaction of 2 with n-butyl lithium was carried out in a similar manner as described above. After separation, 0.57g(53%) of 2'S,5'S-5'-n-butylnicotine(3c) and 0.19g(18%) of 2'S,5'R-5'-n-butylnicotine(4c) were obtained. (3c): mass(m/z); 218(M+; 1), 92(4), 118(5), 130(6), 161(100), 171(10), 217(1): Found: m/z, 218.1761. Calcd for C₁₄H₂₂N₂: M, 218.1783; [α]_D²⁵ -37.6° (c=1.9, MeOH); (4c): mass(m/z); 218(M+; 2), 118(3), 130(29), 161(100), 162(11), 217(1): Found: m/z, 218.1759. Calcd for C₁₄H₂₂N₂: M, 218.1783; [α]_D²⁵ -85.1° (c=0.1, MeOH).

2'S,5'R-5'-Phylnicotine(3d) and 2'S,5'S-5'-Phylnicotine(4d) --- The reaction of 2 with phenyl lithium was carried out in a similar manner as

described above. After separation, 0.71g(59%) of 2'S,5'R-5'-phenylnicotine(3d) and 0.08g(7%) of 2'S,5'S-5'-phenylnicotine(4d) were obtained. (3d): mass(m/z); 238(M+; 21), 91(8), 118(13), 130(15), 160(96), 161(100), 195(4), 209(15), 237(33): Found: m/z, 238.1493. Calcd for C₁₆H₁₈N₂: M, 238.1470: [α]_D²⁵ -8.3° (c=6.1, MeOH); (4d): mass(m/z); 238(M+; 30), 91(12), 118(16), 130(17), 160(100), 161(79), 195(9), 209(33), 237(30): Found: m/z, 238.1441. Calcd for C₁₆H₁₈N₂: M, 238.1470: [α]_D²⁵ -88.9° (c=1.26, MeOH).

2'S,5'R-5'-Isopropylnicotine(3e) and 2'S,5'S-5'-Isopropylnicotine(4e) --- The reaction of 2 with isopropyl lithium was carried out in a similar manner as described above. After separation, 0.48g(48%) of 2'S,5'R-5'-isopropylnicotine (3e) and 0.06g(6%) of 2'S,5'S-5'-isopropylnicotine(4e) were obtained. (3e): mass(m/z); 204(M+; 1), 118(5), 130(50), 161(100), 162(17), 189(1), 203(1): Found: m/z, 204.1611. Calcd for C₁₃H₂₀N₂: M, 204.1626: [α]_D²⁵ -29.5° (c=0.6, MeOH); (4e): mass(m/z); 204(M+; 1), 118(7), 130(45), 161(100), 162(11), 189(3): Found: m/z, 204.1650. Calcd for C₁₃H₂₀N₂: M, 204.1626: [α]_D²⁵ -15.0° (c=0.08, MeOH).

2'S,5'R-5'-tert-Butylnicotine(3f) --- The reaction of 2 with tert-butyl lithium was carried out in a similar manner as described above. After separation 0.10g(9%) of 2'S,5'R-5'-tert-butylnicotine(3f) was obtained. mass(m/z); 218(M+; 0.02), 130(25), 161(100), 162(9), 203(5): Found: m/z, 203.1547. Calcd for C₁₃H₁₉N₂: M - CH₃, 203.1548: [α]_D²⁵ -76.0° (c=0.5, MeOH).

REFERENCES

- 1) J. I. Seeman, H. V. Secor, C. R. Howe, C. G. Chavdarian, and L. W. Morgan, J. Org. Chem., **48**, 4899 (1983); C. G. Chavdarian and J. I. Seeman, Tetrahedron Lett., **23**, 2519 (1982); R. M. Acheson, M. I. Ferris, and N. M. Sinclair, J. Chem. Soc. Perkin I, 579 (1980).
- 2) W. Hankins and A. Burger, J. Pharm. Sci., **59**, 342 (1970).
- 3) J. F. Whidby and J. I. Seeman, J. Org. Chem., **41**, 1585 (1976); T. P. Pitner, W. B. Edwards, III, R. L. Bassfield, and J. F. Whidby, J. Am. Chem. Soc., **100**, 246 (1978).
- 4) D. K. Dalling, D. M. Grant, J. Am. Chem. Soc., **89**, 6612 (1967).
- 5) R. Lukes and O. Grossmann, Coll. Czech. Chem. Commun., **8**, 533 (1936).

Received, 14th October, 1985