

HETEROCYCLES, Vol. 89, No. 7, 2014, pp. 1620 - 1631. © 2014 The Japan Institute of Heterocyclic Chemistry
Received, 9th April, 2014, Accepted, 2nd May, 2014, Published online, 27th May, 2014
DOI: 10.3987/COM-14-13003

STEREOCHEMICAL ASSIGNMENT OF FOUR DIASTEREOMERS OF 3,4-DIMETHYLPYROGLUTAMIC ACID, A MOIETY OF CALLIPELTIN B

Mari Kikuchi and Hiroyuki Konno*

Department of Biochemical Engineering, Graduate School of Science and Technology, Yamagata University, Yonezawa, Yamagata 992-8510, Japan

*E-mail address: konno@yz.yamagata-u.ac.jp

Abstract – The synthesis and stereochemical assignment of four diastereoisomers of 3,4-dimethylpyroglutamic acid (pDME) (**1**) are described. Stereo-divergent synthesis of four pDMEs (**1**) was achieved starting from Thottathil's synthon and Garner's aldehyde for comparison of synthetic pDMEs (**1**) by ^1H NMR analysis and CD spectra. The stereochemistry of pDME (**1**) in cyclic depsipeptide callipeltin B (**2**) with cytotoxic and anti-HIV activities was confirmed to be $2S,3S,4R$. Coupling constant between H2 and H3 of synthetic ($2S,3S,4R$)-pDME (**1a**) showed a smaller value than those of other isomers. In addition, synthetic pDMEs (**1**) hardly showed cytotoxicity against HeLa cells.

INTRODUCTION

3,4-Dimethylpyroglutamic acid (pDME) (**1**) is an unusual amino acid, including callipeltin B (**2**) isolated from marine sponge *Callipelta* sp.¹ 3,4-Dimethylglutamine (DMQ) (**3**), an acyclic unusual amino acid as a precursor of pDME (**1**), is a common component of callipeltin A (**4**),² papuamide A,³ neamphamides,⁴ theopapuamides,⁵ mirabamides,⁶ homophymines⁷ and pipecolidepsin A.⁸ The stereochemistry of DMQ (**3**) was determined to be $2S,3S,4R$ on the basis of a positive Cotton effect on the CD spectrum and NMR analysis of the hydrolysate, which was pDME (**1**) derived from callipeltin A (**4**). Therefore, the stereochemistry of pDME (**1**) in callipeltin B (**2**) was also concluded to be $2S,3S,4R$. Synthetic studies to construct unusual amino acids in the above-mentioned cyclic depsipeptides have been reported by several groups.⁹⁻¹⁴ In particular, synthesis of pDME (**1a**) is important research to achieve the total synthesis of complexed cyclic depsipeptides since DMQ (**3**) could be converted by a ring-opening reaction for pDME (**1a**).¹² Preparation of DMQ and pDME derivatives has been reported independently by Lipton's and Hamada's groups to date. Lipton *et al.* obtained pDME (**1a**) in 9 steps via cuprate addition and enolate

alkylation, followed by kinetic epimerization of the methyl substituent starting from L-pyroglutamic acid (L-pGlu) (**5**).¹² On the other hand, Hamada's group chose a Thottathil's synthon¹⁵ as the starting material for pDME (**1a**) and produced in 6 steps.¹³ For our research program of the synthesis of callipeltins and homophymines, we reported the preparation of L-N-methylalanine, L-N-methylglutamine, (2*S*,3*R*)-D-β-methoxytyrosine (βMOY), D-allothreonine and pDME (**1a**) for the solid phase synthesis of callipeltin B (**2**) in our previous works.¹⁶⁻¹⁹ However, ¹H NMR spectral data of pDME (**1**), which was obtained on large-scale synthesis based on previous synthetic route starting from Garner's aldehyde, and those of natural product¹ were slightly different. Synthetic pDME (**1a**) derived from Garner's aldehyde was simultaneously recognized as the equilibrium mixture. Because of the formal synthesis of pDME (**1a**) derived from Garner's aldehyde in our previous report,¹⁹ the stereochemistry of pDME (**1a**) has been hardly investigated in detail. In addition, spectral data of 3,4-dimethylpyroglutaminol (**7a**) reported by us,^{19,20} which was the substrate of the final oxidation reaction for pDME (**1a**), were similar to those of Hamada's report.¹³ This result might suggest that the stereochemistry of pDME (**1**) needed reassignment by comparison of all diastereoisomers since Lipton's and Hamada's groups synthesized only the natural form and spectral data for comparison of other diastereoisomers were not indicated. In contrast, Lipton's group reported that callipeltin B (**2**) showed cytotoxicity against HeLa cells with IC₅₀ = 98 μM and desmethoxycallipeltin B, in which substitution of D-Tyr for βMOY indicated IC₅₀ = 128 μM.²¹ The results showed that a methoxy group of βMOY was not needed to show cytotoxicity. Furthermore, we supposed that dimethyl groups of pDME (**1a**) were essential for cytotoxicity in our previous study²² using callipeltin B analogues; however, evaluation of pDMEs (**1**) against the cell lines has never been reported. For this reason, we herein disclose the preparation of four possible diastereoisomers, reconfirmation of





Figure 1. Dimethylpyroglutamic acid (pDME) (**1**), natural form of pDME (**1a**) and DMQ (**3**) and callipeltins B (**2**) and A (**4**).

the stereochemistry of the natural form, and evaluation of cytotoxicity against HeLa cells for synthetic pDMEs (**1**) (Figure 1).


RESULTS AND DISCUSSION

At first, we synthesized pDMEs (**1a,b**) using Thottathil's synthon.¹⁵ Preparation of the requisite dimethyl compound **6** was already described in 7 steps starting from L-pGlu (**5**) by Hamada and co-workers and therefore verification of the stereochemistry of the starting material (**6**) was performed correctly. After the spectral data of the obtained **6** were found to be identical to those of Hamada's group, cleavage of *N,O*-acetal of **6** using 9% TFA/CH₂Cl₂ proceeded smoothly to afford alcohol (**7b**) in quantitative yield. Direct epimerization of H4-proton and/or oxidation of the hydroxy group of **7b** afforded the expected products in extremely low yield. Thereby, after the protection of **7b** with TBS and Boc groups, treatment with NaHMDS at -80 °C gave a mixture of the desired isomerized product (**8a**) and substrate (**8b**) in a ratio of 3:1 in satisfactory yield. Separation of the mixture of **8a** and **8b** afforded pure **8a** by column chromatography in 67% yield, which is a known product reported by Lipton's group.¹² Boc carboxylic acid (**9a**) derived from **8a** was obtained in the Jones condition with CrO₃/H₂SO₄ in acetone at 0 °C (deprotection of TBS group and oxidation) and subsequently, deprotection of the Boc group with 50% TFA/CH₂Cl₂ proceeded to give the desired (2*S*,3*S*,4*R*)-pDME (**1a**) according to Lipton's procedures (Scheme 1).¹² It was noted that treatment of **9a** with neat TFA afforded an undesired product (**1b**) by epimerization of H4-proton. Similarly, (2*S*,3*S*,4*S*)-pDME (**1b**) as a single diastereomer was given from **8b** using the Jones condition followed by cleavage of the Boc group in 68% yield over 2 steps. As a result, ¹H, ¹³C NMR, CD, IR, mp, [α]_D and MS spectral data of synthetic **1a** resembled those of the natural product¹ and Hamada's group,¹³ and furthermore, an equilibrium mixture of synthetic **1a** could hardly be observed. On the contrary, pDME (**1a**) described in our previous report was not identified. Therefore, we



Scheme 1. Synthesis of (2*S*,3*S*,4*R*)- and (2*S*,3*S*,4*S*)-pDME (**1a,b**).


synthesized pDME (**1a**) according to the previous route to reassign the spectral data of all compounds obtained from previous study.¹⁹ As a result, we focused on the reaction of diastereoselective hydrogenation of **10**. Thereby, hydrogenation of olefin (**10**), which was obtained starting from serine derivatives in 7 steps, by H₂/Pd-C in MeOH/AcOEt afforded a single product in quantitative yield. In contrast, Hanessian has demonstrated that treatment of olefin (**11**) with Gilman reagent gave ethyl ester (**12**) in 98% yield at the high diastereoselectivity by internal 1,2-asymmetric induction.²³⁻²⁶ ¹H NMR spectral data of **12** derived from **11** by Hanessian's protocol were clearly identical to the product obtained by hydrogenation of **10** in the present study (Scheme 2). Hanessian's 1,4-addition reaction with Me₂CuLi/TMSCl for **12** was proposed via Felkin-Ahn type conformation.²⁷ We conjectured that **13** would be obtained due to the addition of hydrogen from the bottom at hydrogenation via similar conformation. However, both spectral data of **12** by Hanessian and our synthetic **13** in the previous literature¹⁹ conformed completely. In other words, two plausible reactions in Hanessian's and our groups proceeded via distinct mechanisms to give the same compound (**12**). Kim *et al.* reported stereoselective dihydroxylation reaction of γ -amino- α,β -unsaturated esters via *N*-outside or *N*-inside conformers.^{28,29} According to these results, we assumed that the hydrogenation of *E*-olefin (**10**) proceeded via *N*-inside conformer to give **12** (Scheme 2).



Scheme 2. Diastereoselection by internal 1,2-asymmetric induction.

The improved synthesis of other pDME isomers (**1c,d**) was carried out for comparison of spectral data. **12** was treated with NaHMDS and MeI in the presence of HMPA to provide the dimethylated product (**14**) in 92% yield as a single isomer. In the process of the purification of **14**, pyrrolidinone (**7c**), transformed via deprotection of *N, O*-acetal, was afforded in trace amount. Therefore, treatment of **14** with SiO₂ in toluene at reflux proceeded pyrrolidinone formation to afford **7c** and its isomer (**7d**) in 54% yield at a ratio of 5:1 with amino lactone (**15**) as a byproduct. In an attempt to separate the mixture of **7c** and **7d**, we found that

these isomers (**7c,d**) easily isomerized under neutral conditions at room temperature for a few hours. Spectral data of the mixture of **7c** and **7d** resembled those of **7a**²⁰ and **7b**, thus determination of the natural form was extremely difficult. In our previous study, we concluded that alcohol (**7**) existed as a rotamer in the ¹H NMR spectrum. Consequently, it was not easy to determine the natural form using **7a-d**, although we have reported the synthesis of pDME (**1a**). Since TEMPO mediated oxidation using the mixture of **7c** and **7d** gave **1c** and **1d** in 77% yield at a ratio of 3:1, their coupling constants between H2 and H3-proton were indicated as $J_{H_2, H_3} = 7.8\sim 8.7$ Hz, which did not conform with those of the natural compound and synthetic **1a** or **1b**. These results propose that the stereochemistry of **1c** and **1d** is *2S,3R,4R* and *2S,3R,4S* (Scheme 3).³⁰ By measurement of the CD spectrum, **1a** and the mixture of **1c** and **1d** showed a positive Cotton effect at 211 and 217 nm, respectively. These CD spectra were similar to those of the natural product having a positive Cotton effect at 208 nm. Because of the negative Cotton effect at 219 nm, **1b** could be rejected completely.



Scheme 3. Synthesis of (*2S,3R,4R*)- and (*2S,3R,4S*)-pDME (**1c,d**).

Next, the molecular force field for four diastereoisomers of pDME (**1**) was calculated by SPARTAN. Four stable conformers were consequently indicated for each diastereoisomer and the energies of most stable conformations were calculated as -29.16, -30.09, -21.13 and -20.44 kJ/mol for **1a-d**, respectively. On account of these results, the dihedrals of H₃C₃C₂H₂ of **1a-d** were 93.5°, 153.6°, 30.8° and 35.7°. These coupling constants calculated by the Karplus equation confirmed that the observed value of the H2-proton of **1a** was much smaller than other forms (Figure 2). We concluded that the stereochemistry of natural pDME (**1**) was *2S,3S,4R* through the synthesis of four diastereoisomers and its simulation. (*2S,3S,4R*)-**1a** only indicated a small coupling constant between H2 and H3-protons at $J_{H_2, H_3} = 3.2$ Hz, and the other three isomers (**1b-d**) had similar coupling constants of $J_{H_2, H_3} = 7.8\sim 8.7$ Hz. These results were confirmed by the comparison of ¹H NMR spectral data of natural callipeltin B (**2**) ($J_{H_2, H_3} = 3.0$ Hz). In addition, **1a** and **1b** have never been observed in the equilibrium state in spite of the isomerized mixture of **1c** and **1d** with H4-proton in neutral conditions.









				
Molecular Mechanics				
	-29.16 kJ/mol	-30.09 kJ/mol	-21.13 kJ/mol	-20.44 kJ/mol
Calcd. dihedral	$\phi = 93.5^\circ$	$\phi = 153.6^\circ$	$\phi = 30.8^\circ$	$\phi = 35.7^\circ$
Calcd. J_{H_2, H_3} (Hz)	2.5	9.8	7.3	6.8
Experimental J_{H_2, H_3} (Hz)	3.2	8.2	8.7	7.8

Figure 2. Stable conformation of pDMEs (**1a-d**) by SPARTAN.

In our previous works on the cytotoxicity of callipeltin B analogues, we suggested that dimethyl groups of pDME (**1**) might be important for potent activities.²² A cytotoxicity assay was performed against HeLa cells using dimethyl and normal pyroglutamic acids, **1a-d**, L-pGlu (**5**), D-pGlu (ent-**5**) for cytotoxicity alongside rotenone, which served as the control. The cytotoxicities of these compounds were determined by measuring live-cell succinate-tetrazolium reductase activity (MTT assay). The normal L- and D-pGlu (**5**, ent-**5**), pDMEs (**1a, b**), and the mixture of **1c** and **1d** showed practically no cytotoxicity. The results suggested that dimethyl groups of pDME (**1**) needed to be incorporated into the peptide, callipeltin B (**2**), for cytotoxicity against HeLa cells.

In conclusion, we prepared four diastereoisomers of pDMEs (**1a-d**) for the first time, and reconfirmation the stereochemistry of natural pDME (**1**) in callipeltin B (**2**) to be *2S,3S,4R*. We supposed that **10** gave **12** by hydrogenation in high diastereoselectivity via *N*-inside conformer. Furthermore, ¹H NMR spectral data of **7a-d** were similar and consequently it was difficult to identify the stereochemistry of natural form pDME (**1**). However the comparison of all diastereomers of pDME (**1**) was efficient, especially using ¹H NMR analysis and CD spectra. We revise the assignment of the stereochemistry of synthetic compounds in our previous research.¹⁹ In addition, cytotoxicity against HeLa cells was shown by the peptide with pDME (**1**). A synthetic study of callipeltin B (**2**) containing synthetic (*2S,3S,4R*)-pDME (**1a**) on a solid support is now underway.

EXPERIMENTAL

General. All solvents were reagent grade. CH₂Cl₂ was distilled from CaH₂. All commercial reagents were of the highest purity available. Analytical TLC was performed on silica gel (Merck Silica gel 60 F₂₅₄). Column chromatography was carried out on Silica Gel 60 N [Kanto, particle size, (spherical, neutral) 40-100 μm or 63-210 μm]. ¹H (400 MHz) and ¹³C NMR (100, 75 MHz) spectra were recorded on either a JEOL JNM-ECX400 or a BRUKER AM-300. Chemical shifts are expressed in ppm relative to TMS (0

ppm), CHCl_3 (7.26 ppm for ^1H and 77.1 ppm for ^{13}C) or H_2O (4.79 ppm for ^1H). IR spectra were obtained on a HORIBA FT-720 spectrometer. Optical rotations were recorded on a JASCO DIP-371 polarimeter at the sodium D line. High-resolution mass spectra (HRMS) were obtained using a JEOL AccuTOF JMS-T100LC (ESIMS) or a JMS-T100GC-EI (EIMS). Melting points were determined on an AS ONE ATM-02. Conformational searches and stable energy calculations were performed with Spartan'14 Parallel Suite for Windows (Wavefunction, Inc). Absorbances for cytotoxicity assay were measured with the CORONA MTP-310Lab. The CD spectra were recorded on a JASCO J-820.

(3*S*,4*S*,5*S*)-5-Hydroxymethyl-3,4-dimethylpyrrolidin-2-one (7b). To a solution of **6** (235 mg, 1.02 mmol) in CH_2Cl_2 (6 mL) was added TFA (600 μL). After the stirring for 1.5 h at room temperature, the mixture was evaporated under reduced pressure. The product was purified with silica gel column chromatography ($\text{CHCl}_3/\text{MeOH}=95:5$) to give **7b** (146 mg, 1.02 mmol, quant.) as a brown oil. $[\alpha]_{\text{D}}^{27}$ 45.8 (*c* 1.0, CHCl_3); ^1H NMR (CDCl_3 , 400 MHz) δ : 1.13 (3H, d, $J=6.4$ Hz), 1.14 (3H, d, $J=7.2$ Hz), 1.63 (1H, ddq, $J=10.4, 8.4, 6.8$ Hz), 2.08 (1H, dq, $J=10.4, 7.2$ Hz), 3.31 (1H, td, $J=7.6, 2.4$ Hz), 3.46 (1H, dd, $J=11.6, 7.6$ Hz), 3.76 (1H, d, $J=11.6$ Hz); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 13.9, 16.5, 40.0, 44.6, 62.4, 64.2, 180.6; IR ν_{max} (film) cm^{-1} : 3338, 2962, 2931, 2875, 1685; EIHRMS m/z $[\text{M}]^+$: 143.0909, Calcd. for $\text{C}_7\text{H}_{13}\text{NO}_2$: 143.0946.

(3*S*,4*S*,5*S*)-*N*-(*tert*-Butoxycarbonyl)-5-(*tert*-butyldimethylsilyloxy)methyl-3,4-dimethylpyrrolidin-2-one (8b). To a solution of **7b** (288 mg, 2.01 mmol) in DMF (6 mL) was added TBSCl (603 mg, 4.02 mmol) and imidazole (545 mg, 8.04 mmol) and then the mixture was stirred at ~ 50 $^\circ\text{C}$. After being stirred for 3.5 h, H_2O and AcOEt were added to the solution. The organic layer was washed with brine, dried over MgSO_4 , and evaporated under reduced pressure. The residue was purified with silica gel column chromatography ($\text{CHCl}_3/\text{MeOH}=99:1$) to give TBS ether (453 mg, 1.76 mmol, 88%) as a pale yellow oil. $[\alpha]_{\text{D}}^{28}$ 31.8 (*c* 0.4, CHCl_3); ^1H NMR (CDCl_3 , 400 MHz) δ : 0.05 (5H, s), 0.08 (1H, s), 0.87 (8H, s), 0.90 (1H, s), 1.13 (3H, d, $J=7.2$ Hz), 1.16 (3H, d, $J=6.8$ Hz), 1.55 (1H, m), 2.05 (1H, m), 3.27 (1H, td, $J=8.2, 2.8$ Hz), 3.38 (1H, dd, $J=9.6, 8.7$ Hz), 3.74 (1H, dd, $J=10.1, 2.8$ Hz), 5.87 (1H, brs); ^{13}C NMR (CDCl_3 , 100 MHz) δ : -5.4, 14.2, 16.9, 18.2, 25.8, 40.6, 44.4, 61.4, 66.0, 179.2; IR ν_{max} (film) cm^{-1} : 3218, 3106, 2956, 2929, 2857, 1704, 1122, 839; ESIHRMS m/z $[\text{M}+\text{Na}]^+$: 280.1708, Calcd. for $\text{C}_{13}\text{H}_{27}\text{NNaO}_2\text{Si}$: 280.1709. To a solution of TBS ether (43.8 mg, 170 μmol) in MeCN (1 mL) was added $(\text{Boc})_2\text{O}$ (371 mg, 1.70 mmol) and DMAP (cat.). After the stirring for 1.5 h at room temperature, H_2O and AcOEt were added to the solution. The organic layer was washed with brine, dried over MgSO_4 , and evaporated under reduced pressure. The residue was purified with silica gel column chromatography (hexane/AcOEt=6:1) to give **8b** (58.9 mg, 165 μmol , 97%) as a pale yellow solid. mp 50-52 $^\circ\text{C}$; $[\alpha]_{\text{D}}^{26}$ -55.5 (*c* 2.7, CHCl_3); ^1H

NMR (CDCl₃, 400 MHz) δ : 0.03 (3H, s), 0.04 (3H, s), 0.87 (9H, s), 1.15 (3H, d, $J=6.9$ Hz), 1.22 (3H, d, $J=7.4$ Hz), 1.53 (9H, s), 1.96 (1H, m), 2.09 (1H, m), 3.54 (1H, td, $J=4.1, 2.3$ Hz), 3.69 (1H, dd, $J=10.5, 1.8$ Hz), 4.03 (1H, dd, $J=10.6, 4.1$ Hz); ¹³C NMR (CDCl₃, 100 MHz) δ : -5.4, 14.8, 18.3, 18.6, 25.8, 28.1, 35.0, 45.4, 61.3, 64.4, 82.6, 150.5, 176.6; IR ν_{\max} (film) cm⁻¹: 2958, 2931, 2857, 1789, 1754, 1716, 1340, 1201, 837; ESIHRMS m/z [M+H]⁺: 358.2451, Calcd. for C₁₈H₃₆NO₄Si: 358.2414.

(3R,4S,5S)-N-(tert-Butoxycarbonyl)-5-(tert-butyldimethylsilyloxy)methyl-3,4-dimethylpyrrolidin-2-one (8a). To a solution of **8b** (1.17 g, 3.28 mmol) in THF (15 mL) at -80 °C was added NaHMDS in THF (2.30 mL, 1.9 mol/L, 4.26 mmol) and then the mixture was stirred at -80 °C. After being stirred for 1 h, AcOH (377 μ L) was added. The mixture was quenched by saturated NH₄Cl *aq.* and diluted with Et₂O. The solution was allowed to warm to room temperature and then extracted with Et₂O. The organic layer was washed with brine, dried over MgSO₄, and evaporated under reduced pressure. The residue was purified with silica gel column chromatography (hexane/AcOEt=8:1) to give **8a** (786 mg, 2.20 mmol, 67%) as a pale yellow oil. $[\alpha]_D^{28}$ -43.5 (c 1.7, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ : 0.03 (3H, s), 0.05 (3H, s), 0.87 (9H, s), 0.99 (3H, d, $J=7.2$ Hz), 1.08 (3H, d, $J=8.0$ Hz), 1.54 (9H, s), 2.44 (1H, m), 2.97 (1H, m), 3.67 (1H, td, $J=5.2, 2.8$ Hz), 3.75 (1H, dd, $J=10.8, 2.8$ Hz), 3.85 (1H, dd, $J=10.8, 5.2$ Hz); ¹³C NMR (CDCl₃, 100 MHz) δ : -5.6, 10.1, 15.7, 18.1, 25.8, 28.0, 32.9, 40.6, 63.0, 64.7, 82.6, 150.5, 176.7; IR ν_{\max} (film) cm⁻¹: 2956, 2931, 2857, 1789, 1754, 1712, 1309, 1159, 837; ESIHRMS m/z [M+Na]⁺: 380.2212, Calcd. for C₁₈H₃₅NNaO₄Si: 380.2233.

(2S,3S,4R)-N-(tert-Butoxycarbonyl)-3,4-dimethylpyroglutamic acid (9a). To a solution of **8a** (317 mg, 887 μ mol) in acetone (5 mL) at 0 °C was added Jones reagent (30% CrO₃/H₂SO₄, 225 μ L). After being stirred for 6.5 h at 0 °C, the reaction was quenched by addition of 2-propanol (225 μ L) at 0 °C and saturated NaHCO₃ *aq.* (125 μ L) was added. The mixture was diluted with H₂O and extracted with AcOEt. The aqueous layers were combined and acidified with 0.1 M HCl. The solution was extracted with AcOEt. The combined organic layers were washed with brine, dried over MgSO₄, and evaporated under reduced pressure. The residue was purified with silica gel column chromatography (CHCl₃/MeOH=98:2 to 95:5) to give **9a** (199 mg, 773 μ mol, 87%) as a colorless amorphous. mp 112-113 °C; $[\alpha]_D^{29}$ -8.0 (c 0.6, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ : 1.12 (3H, d, $J=7.3$ Hz), 1.13 (3H, d, $J=7.3$ Hz), 1.52 (9H, s), 2.58 (1H, m), 2.82 (1H, m), 4.24 (1H, d, $J=1.8$ Hz); ¹³C NMR (CDCl₃, 100 MHz) δ : 9.9, 15.3, 27.9, 34.3, 40.7, 64.2, 83.9, 149.9, 175.7; IR ν_{\max} (film) cm⁻¹: 3218, 2979, 2937, 2879, 1781, 1751, 1720, 1311, 1155; ESIHRMS m/z [M+Na]⁺: 280.1137, Calcd. for C₁₂H₁₉NNaO₅: 280.1161.

(2S,3S,4R)-3,4-Dimethylpyroglutamic acid (1a). To a solution of **9a** (49.5 mg, 192 μ mol) in CH₂Cl₂

(2 mL) was added TFA (2 mL). After the stirring for 1.5 h at room temperature, the mixture was evaporated under reduced pressure to give **1a** (30.2 mg, 192 μmol , quant.) as a white powder. mp 177-180 °C; $[\alpha]_{\text{D}}^{29}$ 48.3 (*c* 0.3, MeOH); CD $[\theta]_{211\text{nm}}$ +29728 (*c* 0.001 M, H₂O); ¹H NMR (D₂O, 400 MHz) δ : 1.05 (3H, d, *J*=6.8 Hz), 1.13 (3H, d, *J*=6.8 Hz), 2.68 (2H, m), 3.96 (1H, d, *J*=3.2 Hz); ¹³C NMR (D₂O, 100 MHz) δ : 9.5, 13.7, 38.0, 39.2, 61.4, 176.3, 183.8; IR ν_{max} (film) cm^{-1} : 3259, 2971, 1735, 1720, 1155; EIHRMS *m/z* [M]⁺: 157.0749, Calcd. for C₇H₁₁NO₃: 157.0739.

(2S,3S,4S)-N-(tert-Butoxycarbonyl)-3,4-dimethylpyroglutamic acid (9b). To a solution of **8b** (137 mg, 530 μmol) in acetone (2.5 mL) at 0 °C was added Jones reagent (30% CrO₃/H₂SO₄, 100 μL). After being stirred for 3.5 h at 0 °C, the reaction was quenched by addition of 2-propanol (100 μL) at 0 °C and saturated NaHCO₃ *aq.* (50 μL) was added. The mixture was diluted with H₂O and extracted with AcOEt. The aqueous layers were combined and acidified with 0.1 M HCl. The solution was extracted with AcOEt. The combined organic layers were washed with brine, dried over MgSO₄, and evaporated under reduced pressure. The residue was purified with silica gel column chromatography (CHCl₃/MeOH=99:1 to 80:20) to give **9b** (73.9 mg, 287 μmol , 76%) as a pale yellow oil. $[\alpha]_{\text{D}}^{23}$ -29.9 (*c* 0.7, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ : 1.24 (3H, d, *J*=7.3 Hz), 1.30 (3H, d, *J*=6.9 Hz), 1.50 (9H, s), 1.94 (1H, m), 2.21 (1H, m), 4.07 (1H, d, *J*=8.2 Hz); ¹³C NMR (CDCl₃, 100 MHz) δ : 13.9, 17.2, 27.9, 38.8, 45.6, 64.4, 84.1, 149.4, 175.1; IR ν_{max} (film) cm^{-1} : 3219, 2978, 2937, 1778, 1753, 1724, 1155; ESIHRMS *m/z* [M+Na]⁺: 280.1175, Calcd. for C₁₂H₁₉NNaO₅: 280.1161.

(2S,3S,4S)-3,4-Dimethylpyroglutamic acid (1b). To a solution of **9b** (65.9 mg, 256 μmol) in CH₂Cl₂ (800 μL) was added TFA (800 μL). After the stirring for 2 h at room temperature, the mixture was evaporated under reduced pressure to give **1b** (36.1 mg, 230 μmol , 90%) as a colorless amorphous. mp 129-130 °C; $[\alpha]_{\text{D}}^{24}$ -13.3 (*c* 0.4, MeOH); CD $[\theta]_{219\text{nm}}$ -26424 (*c* 0.001 M, H₂O); ¹H NMR (D₂O, 400 MHz) δ : 1.06 (3H, d, *J*=7.3 Hz), 1.22 (3H, d, *J*=6.9 Hz), 2.03 (1H, m), 2.19 (1H, m), 3.88 (1H, d, *J*=8.2 Hz); ¹³C NMR (D₂O, 100 MHz) δ : 13.5, 16.6, 42.7, 43.9, 61.2, 175.9, 182.9; IR ν_{max} (film) cm^{-1} : 3276, 2969, 2929, 1739, 1722, 1660; EIHRMS *m/z* [M]⁺: 157.0752, Calcd. for C₇H₁₁NO₃: 157.0739.

(3R,4R,5S)-5-Hydroxymethyl-3,4-dimethylpyrrolidin-2-one (7c) and (3S,4R,5S)-5-hydroxymethyl-3,4-dimethylpyrrolidin-2-one (7d). To a solution of **14** (195 mg, 647 μmol) in toluene (10 mL) was added SiO₂ (2.0 g) and heated to reflux for 8 h. The mixture was filtrated, washed with 30% MeOH/CHCl₃ and evaporated under reduced pressure. The residue was purified with silica gel column chromatography (CHCl₃/MeOH=97:3 to 95:5) to give mixture of **7c** and **7d** (5:1, 50.4 mg, 352 μmol , 54%) as a brown oil. ¹³C NMR (CDCl₃, 75 MHz) δ : 13.1, 13.8, 13.9, 16.4, 39.8, 40.0, 42.6, 44.4, 57.3,

58.4, 62.0, 62.1, 63.9, 180.4, 181.6; IR ν_{\max} (film) cm^{-1} : 3385, 2962, 2931, 2877, 1685; EIHRMS m/z $[\text{M}]^+$: 143.0944, Calcd. for $\text{C}_7\text{H}_{13}\text{NO}_2$: 143.0946; **7c**: ^1H NMR (CDCl_3 , 400 MHz) δ : 1.11-1.14 (6H, m), 2.16 (2H, m), 3.55-3.76 (3H, m). **7d**: ^1H NMR (CDCl_3 , 400 MHz) δ : 1.13-1.15 (6H, m), 1.69-2.04 (2H, m), 3.31 (1H, m), 3.45 (1H, m), 3.79 (1H, m).

(2S,3R,4R)-3,4-Dimethylpyroglutamic acid (1c) and (2S,3R,4S)-3,4-dimethylpyroglutamic acid (1d).

To a solution of mixture **7c** and **7d** (105 mg, 732 μmol) in MeCN (500 μL) and 0.5 M phosphate buffer (pH 6.6) (500 μL) was added NaClO (2 mL), NaClO₂ (132 mg, 1.46 mmol) and TEMPO (16.3 mg, 105 μmol). After the stirring for 4.5 h at room temperature, 1 M HCl and AcOEt were added to the solution. The organic layer was washed with brine, dried over MgSO₄, and evaporated under reduced pressure. The residue was purified with silica gel column chromatography ($\text{CHCl}_3/\text{MeOH}=95:5$) to give mixture of **1c** and **1d** (3:1, 88.8 mg, 565 μmol , 77%) as a colorless oil. CD $[\theta]_{217\text{nm}} +21097$ (c 0.001 M, H₂O); ^{13}C NMR (D_2O , 100 MHz) δ : 14.3, 14.4, 14.6, 15.0, 41.2, 41.6, 43.0, 43.2, 69.2, 71.0, 173.9, 179.8, 180.0, 181.3; IR ν_{\max} (film) cm^{-1} : 3417, 2972, 2939, 1732, 1716, 1213; EIHRMS m/z $[\text{M}]^+$: 157.0730, Calcd. for $\text{C}_7\text{H}_{11}\text{NO}_3$: 157.0739; **1c**: ^1H NMR (D_2O , 400 MHz) δ : 1.07 (3H, d, $J=7.7$ Hz), 1.11 (3H, d, $J=7.2$ Hz), 2.27 (1H, m), 2.47 (1H, m), 4.37 (1H, d, $J=8.7$ Hz). **1d**: ^1H NMR (D_2O , 400 MHz) δ : 1.14 (3H, d, $J=6.9$ Hz), 1.22 (3H, d, $J=6.9$ Hz), 2.11 (1H, m), 2.28 (1H, m), 4.04 (1H, d, $J=7.8$ Hz).

Chemical shift calculations. Synthetic pDMEs (**1a-d**) were submitted to a conformational search employing Molecular Mechanics to afford four stable conformers, respectively. The most stable conformer about each diastereomer was calculated the energies, dihedrals of $\text{H}_3\text{C}_3\text{C}_2\text{H}_2$ and chemical shifts.

Evaluation of the cytotoxicity by MTT assay. HeLa cells were maintained in a suspension culture of DMEM supplemented with 5% FBS (Fetal Bovine Serum) containing 1% of a penicillin-streptomycin mixture. A 100 μL aliquot of HeLa cells (5000 cells/mL) was added to a 96 well plate and incubated for 24 h at 37 °C in a humidified incubator containing 5% CO₂ in air. After 24 h, a 10 μL aliquot of compound **1a**, **1b**, mixture of **1c** and **1d**, L-pGlu (**5**), D-pGlu (ent-**5**) (concentrations varying in the range of 5, 4, 3, 2, 1, 0.1 mM) and rotenone as a control was added to each of the 96 wells and incubated for 24 h. A 10 μL WST-8 solution (mixture of WST-8 and 1-Methoxy PMS) was added to each well and the incubation continued for 2 h. The visible absorbance at 450 nm and 630 nm as the reference wavelength of each well was quantified using a microplate reader.

ACKNOWLEDGEMENTS

This work was supported in part by a Grant-in aid from the Japan Society for the Promotion of the

Science (25450145) to H. K. and M. K. is grateful to the Japan Society for the Promotion of Science Fellowships for young scientists (8696).


SUPPORTING INFORMATION AVAILABLE

¹H, ¹³C NMR, IR and MS spectra of **1a-d**, and **7a-d** and CD spectral data for **1a-d**. This material associated with this article can be found in the online version.

REFERENCES AND NOTES

1. M. V. D'Auria, A. Zampella, L. Gomez-Paloma, and L. Minale, *Tetrahedron*, 1996, **52**, 9589.
2. A. Zampella, M. V. D'Auria, L. Gomez-Paloma, A. Casapullo, L. Minale, C. Debitus, and Y. Henin, *J. Am. Chem. Soc.*, 1996, **118**, 6202.
3. P. W. Ford, K. R. Gustafson, T. C. McKee, N. Shigematsu, L. K. Maurizi, L. K. Pannell, D. E. Williams, E. D. de Silva, P. Lassota, T. M. Allen, R. Van Soest, R. J. Andersen, and M. R. Boyd, *J. Am. Chem. Soc.*, 1999, **121**, 5899.
4. N. Oku, K. R. Gustafson, L. K. Cartner, J. A. Wilson, N. Shigematsu, S. Hess, L. K. Pannell, M. R. Boyd, and J. B. McMahon, *J. Nat. Prod.*, 2004, **67**, 1407.
5. A. S. Ratnayake, T. S. Bugni, X. Feng, M. K. Harper, J. J. Skalicky, K. A. Mohammed, C. D. Andjelic, L. R. Barrows, and C. M. Ireland, *J. Nat. Prod.*, 2006, **69**, 1582.
6. A. Plaza, E. Gustchina, H. L. Baker, M. Kelly, and C. A. Bewley, *J. Nat. Prod.*, 2007, **70**, 1753.
7. A. Zampella, V. Sepe, P. Luciano, F. Bellotta, M. C. Monti, M. V. D'Auria, T. Jepsen, S. Petek, M.-T. Adeline, O. Laprevote, A.-M. Aubertin, C. Debitus, C. Poupat, and A. Ahond, *J. Org. Chem.*, 2008, **73**, 5319.
8. L. Coello, R. Fernandez, J. F. Reyes, A. Francesch, and M. D. C. Cuevas, *Int. Appl. Pat.*, WO 2010/070078 A1, (24 June 2010).
9. R. Krishnamoorthy, L. D. Vazquez-Serrano, J. A. Turk, J. A. Kowalski, A. G. Benson, N. T. Breaux, and M. A. Lipton, *J. Am. Chem. Soc.*, 2006, **128**, 15392.
10. J. P. Genet, S. Juge, and S. Mallart, *Tetrahedron Lett.*, 1998, **29**, 6765.
11. M. Inoue, N. Shinohara, S. Tanabe, T. Takahashi, K. Okura, H. Itoh, Y. Mizoguchi, M. Iida, N. Lee, and S. Matsuoka, *Nat. Chem.*, 2010, **2**, 280.
12. C. M. Acevedo, E. F. Kogut, and M. A. Lipton, *Tetrahedron*, 2001, **57**, 6353.
13. N. Okamoto, O. Hara, K. Makino, and Y. Hamada, *Tetrahedron: Asymmetry*, 2001, **12**, 1353.
14. S. Calimsiz and M. A. Lipton, *J. Org. Chem.*, 2005, **70**, 6218.
15. J. K. Thottathil, J. L. Moniot, R. H. Mueller, M. K. Y. Wong, and T. P. Kissick, *J. Org. Chem.*, 1986, **51**, 3140.

16. M. Kikuchi, K. Nosaka, K. Akaji, and H. Konno, *Tetrahedron Lett.*, 2011, **52**, 3872.
17. H. Konno, S. Aoyama, K. Nosaka, and K. Akaji, *Synthesis*, 2007, 3666.
18. M. Kikuchi and H. Konno, *Tetrahedron*, 2013, **69**, 7098.
19. H. Konno, Y. Takebayashi, K. Nosaka, and K. Akaji, *Heterocycles*, 2010, **81**, 79.
20. Treatment of **8a** with TBAF followed by TFA gave **7a** in moderate yield to use the comparison with other isomers. $[\alpha]_D^{23}$ 76.2 (*c* 0.2, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ : 1.01 (3H, d, *J*=6.9 Hz), 1.07 (3H, d, *J*=7.8 Hz), 2.25 (1H, m), 2.55 (1H, m), 3.33 (1H, brs), 3.49 (1H, td, *J*=10.1, 6.9 Hz), 3.74 (1H, d, *J*=10.5 Hz); ¹³C NMR (CDCl₃, 100 MHz) δ : 10.7, 13.8, 34.5, 40.0, 62.4, 64.3, 181.9; IR ν_{\max} (film) cm⁻¹: 3319, 2970, 2939, 2875, 1672; EIHRMS *m/z* [M]⁺: 143.0929, Calcd. for C₇H₁₃NO₂: 143.0946.



21. R. Krishnamoorthy, B. L. Richardson, and M. A. Lipton, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 5136.
22. M. Kikuchi, Y. Watanabe, M. Tanaka, K. Akaji, and H. Konno, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 4865.
23. S. Hanessian, Y. Gai, and W. Wang, *Tetrahedron Lett.*, 1996, **37**, 7473.
24. S. Hanessian, W. Wang, and Y. Gai, *Tetrahedron Lett.*, 1996, **37**, 7477.
25. H. Yoda, T. Shirai, T. Katagiri, K. Takabe, K. Kimata, and K. Hosoya, *Chem. Lett.*, 1990, 2037.
26. C. Flamant-Robin, Q. Wang, and A. Sasaki, *Tetrahedron Lett.*, 2001, **42**, 8483.
27. M. T. Reetz and D. Rohrig, *Angew. Chem., Int. Ed. Engl.*, 1989, **28**, 1706.
28. J. Jeon, S.-K. Hong, J. S. Oh, and Y. G. Kim, *J. Org. Chem.*, 2006, **71**, 3310.
29. J. S. Oh, J. Jeon, D. Y. Park, and Y. G. Kim, *Chem. Commun.*, 2005, 770.
30. Isomerization from γ -lactone (**17**) to **16** was not observed during a couple of months at room temperature. However, **7d** gradually converted to **7c** over several days to attain the equilibrium state at a ratio of 5:1.

