

Supporting Information

PHENYLPROPANOIDS FROM THE LEAVES OF *NICOTIANA TABACUM* AND THEIR ANTI-TOBACCO MOSAIC VIRUS ACTIVITIES

**Qin-Peng Shen,¹ Lan Li,^{1,2} Xin-Meg Xu,^{a,b} Chun-Bo Liu,¹ Neng-Jun Xiang,¹
Feng-Mei Zhang,¹ Pei He,¹ Zhi-Hua Liu,¹ Shan-Zhai Shang,¹ Ming-Ming
Miao,^{1,*} Guang-Yu Yang^{1,*}**

¹ Key Laboratory of Tobacco Chemistry of Yunnan Province, Research and Development of Center, China Tobacco Yunnan Industrial Co., Ltd., Kunming 650231, P.R. China; ² School of Pharmaceutical Science & Yunnan Key Laboratory of Pharmacology for Nature Products, Kunming Medical University, Kunming, Yunnan 650500, PR China ; E-mail: miaomm2014@163.com ; ygy1110@163.com

Contents of Supporting Information

No.	Contents:	Pages:
Figure S1-S2	¹³ C NMR and ¹ H NMR spectra of 5-hydroxy-6-(3-hydroxypropanoyl)-isobenzofuran-1(3 <i>H</i>)-one (1)	3-4
Figure S3-S4	¹³ C NMR and ¹ H NMR spectra of 6-(3-hydroxypropanoyl)-5-methylisobenzofuran-1(3 <i>H</i>)-one (2)	5-6

Figure S1 ^{13}C NMR spectrum of 5-hydroxy-6-(3-hydroxypropanoyl)-isobenzofuran-1(3*H*)-one (**1**)




Figure S2 ¹H NMR spectrum of 5-hydroxy-6-(3-hydroxypropanoyl)-isobenzofuran-1(3H)-one (**1**)




Figure S3 ^{13}C NMR a spectrum of 6-(3-hydroxypropanoyl)-5-methylisobenzofuran-1(3H)-one (2)




Figure S4 ¹H NMR spectrum of 6-(3-hydroxypropanoyl)-5-methylisobenzofuran-1(3H)-one (2)

