

## **Supporting Information**

### **A PAIR OF NEW NON-GLYCOSIDIC IRIDOID EPIMERS FROM *SCROPHULARIA NINGPOENSIS***



**Wei Zhang, Peng Liu, Xin-Yu Ji, Xiang-Yang Chen, Yan-Yan Jiang, Bin Liu\***

School of Chinese Materia Medica, Beijing University of Chinese Medicine, Beijing  
102488, People's Republic of China

\* Corresponding authors. E-mail: liubinyn67@163.com

## Contents Page

Figure S1	UV spectrum of <b>1</b> in MeOH	3
Figure S2	IR (KBr disc) spectrum of <b>1</b>	3
Figure S3	HR-ESI-MS spectrum of <b>1</b>	4
Figure S4	<sup>1</sup> H NMR spectrum of compound <b>1</b>	4
Figure S5	<sup>13</sup> C NMR spectrum of compound <b>1</b>	5
Figure S6	DEPT-135 spectrum of compound <b>1</b>	5
Figure S7	<sup>1</sup> H- <sup>1</sup> H COSY spectrum of compound <b>1</b>	6
Figure S8	HSQC spectrum of compound <b>1</b>	6
Figure S9	HMBC spectrum of compound <b>1</b>	7
Figure S10	NOESY spectrum of compound <b>1</b>	7



+TOF MS: 0.333 to 0.367 min from Sample 3 (sj7) of 20140624.wiff  
a=3.59928614855269640e-004, t0=-1.72129387158484340e+001 (Turbo Spray) Max. 1.3e4 counts




Figure S3. HR-ESI-MS spectrum of 1







Figure S4.  $^1H$  NMR spectrum of compound 1 in  $CD_3OD$  (500 MHz)




**Figure S5.**  $^{13}\text{C}$  NMR spectrum of compound **1** in  $\text{CD}_3\text{OD}$  (125 MHz)




**Figure S6.** DEPT-135 spectrum of compound **1** in  $\text{CD}_3\text{OD}$  (125 MHz)




**Figure S7.**  $^1\text{H}$ - $^1\text{H}$  COSY spectrum of compound **1** in  $\text{CD}_3\text{OD}$  (500 MHz)



**Figure S8.** HSQC spectrum of compound **1** in  $\text{CD}_3\text{OD}$  (500 MHz)



**Figure S9.** HMBC spectrum of compound **1** in CD<sub>3</sub>OD (500 MHz)



**Figure S10.** NOESY spectrum of compound **1** in CD<sub>3</sub>OD (500 MHz)