

Supporting Information

AN EFFICIENT SYNTHESIS N-ARYL-2-(INDOL-3-YL)ACETAMIDES VIA MULTI-COMPONENT REACTIONS

Guo-Ning Zhang,^a Mengye Zhang,^b Xiaonan Du,^a Mei Zhu,^a Juxian Wang,^{a,*}
Daqing Shi,^{b,*} and Yucheng Wang,^{a,*}

^aInstitute of Medicinal Biotechnology, Chinese Academy of Medical Science and
Peking Union Medical College, Beijing 100050, P. R. China; E-mail:
wangjuxian@imb.pumc.edu.cn (J. W.); wangyucheng@imb.pumc.edu.cn (Y. W.)

^bKey Laboratory of Organic Synthesis of Jiangsu Province, College of Chemistry,
Chemical Engineering and Materials Science, Soochow University, Suzhou 215123,
Jiangsu, P. R. China; E-mail: dqshi@suda.edu.cn


Context

Copies of ¹H NMR and ¹³C NMR Spectra for Compound 4a-4j


S2-S11

Copies of ^1H NMR, ^{13}C NMR and HRMS of compounds


^1H NMR of compound **4a**




^{13}C NMR of compound **4a**




¹H NMR of compound **4b**




¹³C NMR of compound **4b**




¹H NMR of compound **4c**




¹³C NMR of compound **4c**




¹H NMR of compound **4d**




¹³C NMR of compound **4d**




^1H NMR of compound **4e**




^{13}C NMR spectrum of compound **4e**




^1H NMR of compound **4f**




^{13}C NMR of compound **4f**




¹H NMR of compound **4h**




¹³C NMR of compound **4h**




¹H NMR of compound **4i**



¹³C NMR of compound **4i**



¹H NMR of compound **4j**



¹³C NMR of compound **4j**

