

## Supporting Information for

### ON THE ORDER OF ADDITION OF SODIUM DISPERSION IN REDUCTIVE DIBORATIONS OF STILBENE AND 1,2-DIPHENYLCYCLOPROPANE


Shuo Wang, Atsushi Kaga, and Hideki Yorimitsu\*

Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo-ku,  
Kyoto 606-8502, Japan


e-mail: [yori@kuchem.kyoto-u.ac.jp](mailto:yori@kuchem.kyoto-u.ac.jp)

#### Contents

- Figure 1.** <sup>1</sup>H NMR spectrum of **1** obtained by Method 1.
- Figure 2.** <sup>1</sup>H NMR spectrum of **1** obtained by Method 2.
- Figure 3.** <sup>1</sup>H NMR spectrum of **2** obtained by Method 1.
- Figure 4.** <sup>1</sup>H NMR spectrum of **2** obtained by Method 2.



**Figure 1.** <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) spectrum of **1** (*syn/anti* = 93/7) obtained by Method 1.



**Figure 2.** <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) spectrum of **1** (*syn/anti* = 93/7) obtained by Method 2.

