

HETEROCYCLES, Vol. 94, No. 11, 2017, pp. 2081 - 2086. © 2017 The Japan Institute of Heterocyclic Chemistry
 Received, 4th August, 2017, Accepted, 28th September, 2017, Published online, 6th October, 2017
 DOI: 10.3987/COM-17-13790

ULTRASOUND-MEDIATED ONE-POT SYNTHESIS OF THE FULLY FUNCTIONALIZED *N,N'*-(1,3-PROPANE)BISPYRROLES


Huijun Liu,* Qi Ma, Zhen Lu, Qiang Zhao, Jinping Song, Kewei Wang, Feng Feng,* and Yong Guo*

Institute of Applied Chemistry, Institute of Organic Chemistry, College of Chemistry and Environmental Engineering, Shanxi Datong University, Datong, 037009, P.R. China; E-mail: dtdxlhj@163.com

Abstract – Five fully functionalized *N,N'*-(1,3-propane)bispyrroles were synthesized by the *tert*-butyl isocyanide-mediated one-pot reaction of bisimines, aroyl chloride and dimethyl acetylenedicarboxylate under ultrasonic irradiation in good yield. The structures of *N,N'*-(1,3-propane)bispyrroles were characterized by ¹H NMR and ¹³C NMR spectroscopy.


N,N'-(1,3-Propane)bispyrrole and their derivatives has received much attention from the research group due to wide range of biological and materials science applications.¹ As a consequence, much attention has been paid to their preparation by classical methods such as the Knorr reaction,² Paal–Knorr reaction,³ *C–N* cross-coupling reaction,⁴ Michael addition reaction,⁵ Hantzsch pyrrole synthesis,⁶ 1,3-dipolar cycloaddition reaction.⁷

In 2015, we reported a new approach for synthesis of bispyrrole by using the ability of alkynes to undergo the double 1,3-dipolar cycloaddition-extrusion to bis(1,3-oxazolium-5-oxides (Münchnones)).⁸ While bismünchnones are typically prepared in situ from bisglycine derivatives, an inadequate aspect is that can themselves require a multistep synthesis. In 2007, a new isocyanide-mediated reaction of imines and acid chlorides generating the mono-oxazolium salt **1** was discovered by Arndtsen and co-workers.⁹ **1** were trapped by alkynes to form mono-pyrroles **2** (Scheme 1).



Scheme 1. Synthesis of monopyrroles by isocyanide-mediated reaction

As such, we became interested in developing routes to construct bispyrroles. The advantage of ultrasound irradiation to promote organic reactions includes shorter reaction time and high-yields.¹⁰ We describe herein that this assembly of fully functionalized *N,N'*-(1,3-propane)bispyrroles can be obtained via isocyanide-mediated reaction under ultrasonic irradiation.



Scheme 2. Synthesis of *N,N'*-(1,3-propane)bispyrrole derivatives **4a-f**

As shown in Scheme 2, target products **4** were synthesized by the three step one-pot method under ultrasonic irradiation, First, at room temperature, bisimines were treated by aryl chloride for 0.5 h, then, *tert*-butyl isocyanide was added, stirring for 0.5 h; The second, EtN^{*i*}Pr₂ was dropped slowly to the above mixture yield yellow-blue color bis-oxazolium salts **3a**, **3b**, **3d** (when 4-nitrobenzoyl chloride was used, yield violet color bis-oxazolium salts **3c**, **3e**, **3f**); Last, surprisingly, the addition of dimethyl acetylenedicarboxylate (DMAD) to **3** leads to the generation of the corresponding bispyrrole **4** over the course of 2 h (Table 1, entries 3 and 5 for 3 h) under ultrasonic irradiation and the bright and charming colors have vanished. Compound **4f** has not been successfully obtained by this method (Table 1, entry 6).

Table 1. Yield of *N,N'*-(1,3-propane)bispyrrole derivatives **4a-f**

Entry	R	R ¹	Compound	Method A		Method B		Method C ⁷	
				Time (US) / h	Yield (%) ^a	Time / h	Yield (%) ^a	Time / h	Yield (%) ^a
1	H	H	4a	2	72.5	24	59.0	72	30.0
2	H	OMe	4b	2	74.6	24	61.2	-	-
3	H	NO ₂	4c	3	69.0	24	45.3	72	18.0
4	OMe	OMe	4d	2	77.7	24	62.1	-	-
5	OMe	NO ₂	4e	3	62.5	24	46.4	-	-
6	NO ₂	NO ₂	4f	3	0.0	24	0.0	-	-

^a Isolated yields.

We carried out the reaction under conventional conditions (Method B) and compared the results with those obtained under ultrasound conditions (Method A). Thus application of ultrasound not only prevents side products but also reduces reaction time. The yields of products under both ultrasound and room temperature conditions are shown in Table 1. In terms of yield and reaction time, method A is more advantageous than method C (Table 1, entries 1 and 3).

The structures of compounds **4a-e** were determined on the basis of their elemental analyses, ^1H NMR, ^{13}C NMR spectroscopic data. In the ^1H NMR spectra of **4a-e**, the proton of methylene possessed *N* showed triplet at 3.10–3.30 ppm, and it is noteworthy that in the ^1H NMR spectra of all bispyrrole esters, the proton of methyl possessed ester group ($-\text{CO}_2\text{Me}$) showed resonances at 3.6–3.7 ppm, the single peak of **4a** (**4d**) is observed at 3.61 ppm (3.62 ppm), the two single peak of **4b** (**4c**, **4e**) are observed at 3.60 and 3.62 ppm (**4c** 3.62 and 3.63 ppm, **4e** 3.62 and 3.64 ppm), the proton of possessed aryl group showed resonances at > 6 ppm. ^{13}C NMR spectra of compounds **4a-4e** showed resonances at 51–52 ppm (a carbon of methyl of CO_2Me); 114–116 ppm (carbon of pyrrole with methoxycarbonyl group) and 164–165 ppm (carbon of carbonyl group of CO_2Me), both consistent with the presence of the heterocyclic ring.

In conclusion, polysubstituted *N,N'*-(1,3-propane)bispyrroles were successfully synthesized utilizing the *tert*-butyl isocyanide-mediated stepwise fashion one-pot of bisimines, aroyl chloride, dimethyl acetylenedicarboxylate under ultrasonic irradiation in short time, simultaneously providing a modular method to construct these bisheterocycles with facile diversity and high atom economy.

EXPERIMENTAL

Melting points were recorded on a Yanaco hot-stage melting point apparatus and uncorrected. Purification of reaction products was carried out by flash chromatography using EM Reagent silica gel 60 (230-400 mesh). Analytical thin layer chromatography was performed on EM Reagent 0.25 mm silica gel 60-F plates. ^1H NMR spectra were recorded at 400 MHz and ^{13}C NMR spectra were recorded at 100 MHz using a JEOL ECA 400 or ECX 400 spectrometer. Chemical shifts were reported in parts per million (δ) relative to tetramethylsilane (TMS). Data are reported as (ap = apparent, s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, b = broad; coupling constant(s) in Hz; integration). Elemental analyses were performed with a Yanaco MT-5 CHN-Corder. All high resolution mass spectra were obtained on a micro TOF-QII 10203 spectrometer. Ultrasonication was done in a KQ5200DE and a frequency of 40 kHz ultrasound clear with an out power 200 W. All of the organic solvents used in this study were dried over appropriate drying agents.

Starting Materials

Bisimine were prepared by previously reported procedures,¹¹ and used directly without purification. All other chemicals used in this study were commercially available.

Typical Procedure for the Preparation of Compounds 4a-e

Method A: A 25 mL two-neck round bottom flask equipped with a magnetic bar was charged the bisimine (0.1 mmol, 1 equiv.) and MeCN (2 mL), and stirring. Aroyl chloride (0.22 mmol, 2.2 equiv.) was injected by syringe. After 0.5 h at room temperature, *tert*-butyl isocyanide (0.22 mmol, 2.2 equiv.) was injected. After 0.5 h at room temperature, diisopropylethylamine (0.22 mmol, 2.2 equiv.) was injected. After 0.5 h at room temperature, DMAD (0.22 mmol, 2.2 equiv.) was injected and the reaction was carried out under ultrasonic irradiation for 2 or 3 h. The solution was concentrated *in vacuo*, the crude products were purified by silica gel chromatography using CH₂Cl₂/EtOAc acetate (*V* : *V* = 4 : 1) as eluent. The corresponding product was obtained.

Method B: A 25 mL two-neck round bottom flask equipped with a magnetic bar was charged the bisimine (0.1 mmol, 1 equiv.) and MeCN (2 mL), and stirring. Aroyl chloride (0.22 mmol, 2.2 equiv.) was injected by syringe. After 0.5 h at room temperature, *tert*-butyl isocyanide (0.22 mmol, 2.2 equiv.) was injected. After 0.5 h at room temperature, diisopropylethylamine (0.22 mmol, 2.2 equiv.) was injected. After 0.5 h at room temperature, DMAD (0.22 mmol, 2.2 equiv.) was injected and the reaction was carried out at room temperature for 24 h. The solution was concentrated *in vacuo* to give the crude product which was purified by silica gel chromatography using CH₂Cl₂/EtOAc (*V* : *V* = 4 : 1) as eluent. The corresponding product was obtained.

Tetramethyl 1,1'-(propane-1,3-diyl) bis(2,5-diphenyl-1*H*-pyrrole-3,4-dicarboxylate) (4a): Colorless solid, yield: 72.5%. mp 119–121 °C. *R*_f = 0.75. ¹H NMR (CDCl₃, 400 MHz): δ = 1.06–1.12 (m, 2H, CH₂), 3.16–3.20 (t, *J* = 7.6 Hz, 4H, CH₂), 3.61 (s, 12H, CH₃), 7.06–7.09 (d, *J* = 8.4 Hz, 8H, ArH), 7.33–7.42 (m, 12H, ArH). ¹³C NMR (CDCl₃, 100 MHz): δ = 31.5, 41.6, 51.6, 114.7, 128.5, 128.9, 130.2, 130.3, 136.0, 166.1. HRMS (ESI⁺) calcd for C₄₃H₃₈N₂O₈ (MH⁺) 711.7768, found 711.7766. Anal. Calcd for C₄₃H₃₈N₂O₈: C, 72.66; H, 5.39; N, 3.94%. Found: C, 72.43; H, 5.43; N, 3.81.

Tetramethyl 1,1'-(propane-1,3-diyl) bis[2-phenyl-5-(4-methoxy-phenyl)-1*H*-pyrrole-3,4-dicarboxylate] (4b): Colorless solid, yield: 74.6%. mp 82–85 °C, *R*_f = 0.67. ¹H NMR (CDCl₃, 400 MHz): δ = 1.07–1.15 (m, 2H, CH₂), 3.16–3.20 (t, *J* = 8.0 Hz, 4H, CH₂), 3.60 (s, 6H, CH₃), 3.62 (s, 6H, CH₃), 3.87 (s, 6H, CH₃), 6.87–6.89 (d, *J* = 8.8 Hz, 4H, ArH), 7.00–7.02 (d, *J* = 8.8 Hz, 4H, ArH), 7.06–7.08 (d, *J* = 6.8 Hz, 4H, ArH), 7.30–7.42 (m, 6H, ArH). ¹³C NMR (CDCl₃, 100 MHz): δ = 31.6, 41.6, 51.6, 55.3, 113.9, 114.5, 114.6, 122.3, 128.5, 128.8, 130.2, 130.4, 131.5, 135.7, 135.9, 159.9, 165.2, 165.3. HRMS (ESI⁺) calcd for C₄₅H₄₂N₂O₁₀ (MH⁺) 771.8286, found 771.8285. Anal. Calcd for C₄₅H₄₂N₂O₁₀: C, 70.12; H, 5.49; N, 3.63%. Found: C, 69.75; H, 5.57; N, 3.87.

Tetramethyl 1,1'-(propane-1,3-diyl) bis[2-phenyl-5-(4-nitrophenyl)-1*H*-pyrrole-3,4-dicarboxylate] (4c): Yellow solid, yield: 69%. mp 199–201 °C. $R_f = 0.70$. $^1\text{H NMR}$ (CDCl_3 , 400 MHz): $\delta = 0.96\text{--}1.04$ (m, 2H, CH_2), 3.23–3.27 (t, $J = 7.6$ Hz, 4H, CH_2), 3.62 (s, 6H, CH_3), 3.63 (s, 6H, CH_3), 7.12–7.14 (t, $J = 7.6$ Hz, 4H, ArH), 7.24–7.26 (d, $J = 8.8$ Hz, 4H, ArH), 7.37–7.47 (m, 6H, ArH), 8.19–8.21 (d, $J = 8.8$ Hz, 4H, ArH). $^{13}\text{C NMR}$ (CDCl_3 , 100 MHz): $\delta = 29.7, 41.8, 51.9, 53.5, 115.7, 116.0, 123.7, 128.7, 129.4, 129.6, 130.1, 131.2, 133.4, 136.7, 136.8, 147.8, 164.3, 164.8$. HRMS (ESI^+) calcd for $\text{C}_{43}\text{H}_{36}\text{N}_4\text{O}_{12}$ (MH^+) 801.7720, found 801.7716. Anal. Calcd for $\text{C}_{43}\text{H}_{36}\text{N}_4\text{O}_{12}$: C, 64.50; H, 4.53; N, 7.00%. Found: C, 65.09; H, 5.17; N, 6.50.

Tetramethyl 1,1'-(propane-1,3-diyl) bis[2,5-di(4-methoxy-phenyl)-1*H*-pyrrole-3,4-dicarboxylate] (4d): Colorless solid, yield: 77.7%. mp 261–262 °C. $R_f = 0.70$. $^1\text{H NMR}$ (CDCl_3 , 400 MHz): $\delta = 1.11\text{--}1.19$ (m, 2H, CH_2), 3.16–3.20 (t, $J = 7.6$ Hz, 4H, CH_2), 3.62 (s, 12H, CH_3), 3.87 (s, 12H, CH_3), 6.87–6.89 (d, $J = 8.8$ Hz, 8H, ArH), 7.00–7.02 (d, $J = 8.4$ Hz, 4H, ArH). $^{13}\text{C NMR}$ (CDCl_3 , 100 MHz): $\delta = 31.7, 41.5, 51.6, 55.3, 113.9, 114.4, 122.4, 131.5, 135.7, 159.9, 165.4$. HRMS (ESI^+) calcd for $\text{C}_{47}\text{H}_{46}\text{N}_2\text{O}_{12}$ (MH^+) 831.8804, found 831.8800. Anal. Calcd for $\text{C}_{47}\text{H}_{46}\text{N}_2\text{O}_{12}$: C, 67.94; H, 5.58; N, 3.37%. Found: C, 67.94; H, 5.77; N, 3.37.

Tetramethyl 1,1'-(propane-1,3-diyl) bis[2-(4-methoxyphenyl)-5-(4-nitrophenyl)-1*H*-pyrrole-3,4-dicarboxylate] (4e): Yellow solid, yield: 62.5%. mp 98–101 °C. $R_f = 0.55$. $^1\text{H NMR}$ (CDCl_3 , 400 MHz): $\delta = 1.01\text{--}1.09$ (m, 2H, CH_2), 3.22–3.26 (t, $J = 7.6$ Hz, 4H, CH_2), 3.62 (s, 6H, CH_3), 3.64 (s, 6H, CH_3), 3.89 (s, 6H, CH_3), 6.92–6.94 (d, $J = 8.8$ Hz, 4H, ArH), 7.07–7.09 (d, $J = 8.8$ Hz, 4H, ArH), 7.23–7.25 (d, $J = 7.6$ Hz, 4H, ArH), 8.19–8.21 (d, $J = 8.0$ Hz, 4H, ArH). $^{13}\text{C NMR}$ (CDCl_3 , 100 MHz): $\delta = 31.5, 41.8, 51.9, 55.4, 113.9, 114.1, 115.6, 115.7, 121.5, 123.6, 131.2, 131.5, 133.1, 136.7, 136.9, 147.9, 160.3, 164.5, 164.9$. HRMS (ESI^+) calcd for $\text{C}_{45}\text{H}_{40}\text{N}_4\text{O}_{14}$ (MH^+) 862.8317, found 862.8317. Anal. Calcd for $\text{C}_{45}\text{H}_{40}\text{N}_4\text{O}_{14}$: C, 62.79; H, 4.68; N, 6.51%. Found: C, 62.55; H, 5.06; N, 6.15.

ACKNOWLEDGEMENTS

We are grateful to the foundation of the “National Natural Sciences Foundation of China” (No. 21301111), “National Natural Sciences Foundation of Shanxi” (Nos. 2015011017, 2015021049), “Science and Technology Innovation Program Development of Shanxi Higher Education Institutions” (No. 2015181), “Research Project Supported by Shanxi Scholarship Council of China” (No. 21016-105), “Aid project for PhD faculties in the University of Shanxi Datong” (No. 2015-B-03), “Science and Technology Program Development of Datong” (No. 2015019) for financial support.

REFERENCES

- (a) Y. Ding, Y. Tang, W. Zhu, and Y. Xie, *Chem. Soc. Rev.*, 2015, **44**, 1101; (b) W. A. Greenberg,

- E. E. Baird, and P. B. Dervan, *Chem. Eur. J.*, 1998, **4**, 796; (c) K. I. Chane-Ching, J. C. Lacroix, R. Baudry, M. Jouini, S. Aeiyaach, C. Lion, and P. C. Lacaze, *J. Electroanal. Chem.*, 1998, **453**, 139; (d) Y.-H. Chen and J. W. Lown, *J. Am. Chem. Soc.*, 1994, **116**, 3995; (e) J. Hlavatý, V. Papež, and L. Kavan, *Synth. Met.*, 1994, **63**, 209.
- O. A. Attanasi, S. Berretta, L. D. Crescentini, G. Favi, G. Giorgi, F. Mantellini, and S. Nicolini, *Adv. Synth. Catal.*, 2011, **353**, 595.
 - (a) S. Cheraghi, D. Saberi, and A. Heydari, *Catal. Lett.*, 2014, **144**, 1339; (b) V. Polshettiwar and R. S. Varma, *Tetrahedron*, 2010, **66**, 1091; (c) V. Polshettiwar, B. Baruwati, and R. S. Varma, *Chem. Commun.*, 2009, **40**, 1837; (d) R. R. Sagyam, H. Vurimidi, P. R. Padi, and M. R. Ghanta, *J. Heterocycl. Chem.*, 2007, **44**, 923; (e) M. Abid, A. Spaeth, and B. Török, *Adv. Synth. Catal.*, 2006, **348**, 2191; (f) B. K. Banik, S. Samajdar, and I. Banik, *J. Org. Chem.*, 2004, **69**, 213.
 - (a) B. Franck, R. Timmermann, and W. Sirges, *Liebigs Ann. Chem.*, 1993, **10**, 1075; (b) U. Burger and F. Dreier, *Tetrahedron*, 1983, **39**, 2065; (c) U. Burger and F. Dreier, *Helv. Chim. Acta*, 1980, **63**, 1190.
 - H. Meyer, *Liebigs Ann. Chem.*, 1981, **1981**, 1534.
 - L. M. Potikha and V. A. Kovtunencko, *Chem. Heterocycl. Compd.*, 2006, **42**, 741.
 - Z. M. Abdulla, R. P. Iyer, K. G. Akamanchi, M. S. Degani, and E. C. Coutinho, *J. Heterocycl. Chem.*, 2011, **48**, 38.
 - H. Liu, S. Ueta, F. Yagishita, M. Nishiuchi, and Y. Kawamura, *Tetrahedron*, 2015, **71**, 3614.
 - D. J. S. Cyr, N. Martin, and B. A. Arndtsen, *Org. Lett.*, 2007, **9**, 449.
 - J. Lindley and T. J. Mason, *Chem. Soc. Rev.*, 1987, **16**, 275.
 - R. W. Layer, *Chem. Ber.*, 1963, **63**, 489.